

# Short-day Exposure Period For Subterranean Turion Formation in Dioecious Hydrilla

J. N. THAKORE, W. T. HALLER AND D. G. SHILLING<sup>1</sup>

## ABSTRACT

Two experiments conducted in the greenhouse determined that a minimum of 20 short-days (SD) were required for induction of hydrilla [*Hydrilla verticillata* (L.f.) Royle] subterranean turions. Fertilizer was applied to the plants in one study but not the other. There were 15 treatments ranging from 0 to 60 sequential days under SD conditions. Measured parameters included shoot weight, subterranean turion number and weight. The unfertilized plants began producing subterranean turions only after exposure to 20 SD, while in the other study, the fertilized plants produced subterranean turions only after being exposed to 38 SD. The shoot weight of the unfertilized plants decreased with increasing SD exposure; whereas that of the fertilized plants increased. The weight of subterranean turions produced did not significantly differ between the two studies. The unfertilized plants produced 0.17 subterranean turions per day after induction while the fertilized plants produced 0.89 per day.

*Key words:* Photoperiod, subterranean turions, fertilizer, plant nutrients, tubers.

## INTRODUCTION

Hydrilla is an exotic, submersed aquatic macrophyte that has grown aggressively and occurs throughout the Southeastern United States (Langeland 1990). As many as 10 times more subterranean turions (also called tubers) than axillary turions may be produced in a hydrilla infested area (Mitra 1955). Subterranean turions contain one apical meristem capable of regenerating a plant and may remain dormant in the hydrosol for 1 to 5 years (Van and Steward 1990, Yeo et al. 1984). Subterranean turions are the primary reason that hydrilla rapidly regrows after management procedures since subterranean turions are in the hydrosol protected from control methods (Haller 1976).

Subterranean and axillary turion production in the field occurs seasonally in the Southeastern United States under SD conditions from October to April (Netherland 1997, Miller et al. 1993). The critical day length for optimum subterranean turion induction in dioecious hydrilla is 10-12 hours (Haller 1976, Van et al. 1978, McFarland and Barko 1990) and may be regulated by the phytochrome system (Klaine and Ward 1984).

Studies on the effects of temperature, nutrient levels, and plant growth regulators on subterranean turion production have been carried out, but the number of SD necessary to induce subterranean turion production in dioecious hydrilla has not been determined (Netherland 1997). This study was conducted to determine the number of SD required for subterranean turion formation in dioecious hydrilla under two nutrient levels.

## MATERIALS AND METHODS

The study was conducted in two greenhouses, a SD and a long-day (LD) greenhouse, in Gainesville, Florida in the winter of 1994 and repeated in the winter of 1995. The temperatures were maintained at 25 to 30C in the LD greenhouse and between 18 to 22C in the SD greenhouse. The LD photoperiod was 16 hours extended with fluorescent and incandescent lights, with a quantum flux density between 500 and 800  $\mu\text{Em}^{-2}\text{s}^{-1}$  (LI 185 Lambda Quantum Meter). The SD greenhouse received 8 to 10 hours ambient light conditions of 800 to 1000  $\mu\text{Em}^{-2}\text{sec}^{-1}$ .

Subterranean turions were collected from established hydrilla populations grown in outdoor 900 L concrete tanks were washed, and sprouted in distilled water at 22C under LD conditions. Sixteen sprouted turions were planted in 30 by 24 by 12 cm plastic containers filled to the depth of 5 cm with potting soil<sup>2</sup> and sand<sup>3</sup>. Fourteen grams of Osmocote<sup>4</sup> were added to each container in the second study. Sprouted subterranean turions were planted in sixty containers for each study. The plants were placed in 540 L fiberglass tanks (well water) in the LD greenhouse. After a month of growth under LD conditions, plants were transported to the SD greenhouse for exposure to the SD treatments. The treatments ranged from 4 to 60 days under SD conditions and progressed sequentially in increments of 4 days. Every 4 days, 4 containers of hydrilla were transferred from the SD greenhouse back to the LD greenhouse to continue growth. A control group was maintained under LD conditions throughout the study.

A completely randomized design was used for the experiment. Each SD exposure treatment was replicated four times using four containers. After the plants were transferred back to the LD greenhouse, they were allowed to grow for 60 days before harvesting. After 60 LD, shoots were severed at the hydrosol, washed vigorously to remove all soil particles, and

<sup>1</sup>Graduate Research Assistant and Professors, respectively. Department of Agronomy, University of Florida, Gainesville, FL. Received for publication April 26, 1997 and in revised form June 10, 1997. Florida Agricultural Experiment Station, Journal Series No.R-05803.

<sup>2</sup>Southland topsoil, Southern Importers, Inc., P.O. Box 8519, Greensboro, NC. 27419.

<sup>3</sup>Playsand, Readymix Cement, Inc. Contents: Crystalline silica.

<sup>4</sup>Osmocote, 14% total Nitrogen, 14% P<sub>2</sub>O<sub>5</sub>, 14% K<sub>2</sub>O.

dried for 2 weeks at 60C. The soil was also washed thoroughly, recognizable subterranean turions attached to rhizome tips and detached from rhizomes were collected, counted, and dried for two weeks at 60C. Shoot and turion dry weights were determined.

Analysis of variance (ANOVA) was carried out to measure the difference in treatment effects between the two studies. Since the treatment effects of study 1 (unfertilized) were significantly ( $p < 0.05$ ) different from the treatment effects of study 2 (fertilized), the studies were not pooled and statistical analysis was carried out separately for each study. For each study, one-way analysis of variance was used to measure the effects of SD conditions on the dependant variables—shoot weight, subterranean turion number, and subterranean turion weight per container—and orthogonal contrasts were performed for each study by grouping the treatment means. Treatments that caused the formation of subterranean turions were designated as inductive and those that did not cause formation were designated as non-inductive treatments. Orthogonal contrasts were carried out to measure the difference between the two groups. Simple linear regression was conducted for analysis of shoot biomass and non-linear regression of subterranean turion number.

## RESULTS AND DISCUSSION

There was a significant year by treatment interaction ( $p < 0.05$ ) for the shoot weight and subterranean turion number; therefore, the studies will be discussed separately.

### STUDY 1 (UNFERTILIZED)

There were significant differences in shoot biomass, subterranean turion number, and subterranean turion biomass resulting from the length of exposure to SD conditions ( $p < 0.05$ ). The treatment means were pooled into two groups and analyzed to determine at what point the SD conditions cause significantly different outcomes in the shoot biomass, and subterranean number and biomass. The first 20 days of SD were grouped together because there was no subterranean turion production during this interval. This grouping was contrasted with the grouping of all means after 20 SD. The contrast indicated that after 20 SD, the shoot weight, turion number, and turion biomass of hydrilla were significantly ( $p < 0.05$ ) different from the first part of the study. The estimated extent of the differences between the first 20 SD and the rest are presented in Table 1.

The shoot biomass (per container) was greater during the first 20 days than during the rest of the study. This observed decrease in shoot biomass has been found to be characteristic of hydrilla in nature during the winter months (MacDonald 1994). This decline in shoot biomass may be due to the onset of subterranean turion production at day 20 when the plants began allocating carbohydrates to the subterranean turions rather than above-ground growth (Figure 1).

Regression analysis was carried out to further examine the change in shoot weight with increasing SD conditions. The regression line describes the decrease in shoot biomass due to increasing SD conditions.

Subterranean turion number and biomass were significantly higher during the second part of the study than the

TABLE 1. COMPARISON OF POOLED NON-INDUCTIVE AND INDUCTIVE MEANS FOR SUBTERRANEAN TURION NUMBER, SHOOT AND SUBTERRANEAN TURION BIOMASS PER CONTAINER IN STUDY 1 (UNFERTILIZED) AND STUDY 2 (FERTILIZED).

Plant component	$H_0: \mu_{\text{Preinduction}} \text{ vs. } \mu_{\text{Postinduction}}$	Standard Error of differences	p-value
Study 1			
Shoot biomass	7.2 vs. 4.65	0.569	0.0001**
Turion number	0 vs. 3.25	0.63	0.0001**
Turion biomass	0 vs. 3.12	0.12	0.0034*
Study 2			
Shoot biomass	11.2 vs. 18.7	1.08	0.0001**
Turion number	0.3 vs. 11.8	0.72	0.0001**
Turion biomass	0.03 vs. 1.35	0.36	0.0047*

preinduction period with an average of 3.25 turions produced after induction. Since subterranean turions were not produced until after 20 SD, non-linear regression was carried out for further analysis. The linear plateau and the step function models were examined to determine which non-linear model better explains how SD effected subterranean turion production in this study. The linear plateau model assumes




Figure 1. The influence of SD exposure period on the shoot biomass (dry wt/container) of unfertilized and fertilized hydrilla.

that there is a linear increase in subterranean turion production once it has been induced by a given number of SD. Whereas the step function model assumes that there is a threshold effect due to SDs and after induction the number of subterranean turions produced remain the same for all inductive treatments. Statistical analyses using both models demonstrated that the linear plateau model better explains the observed data. Also, the biological effects due to SD did not resemble a threshold type response modeled by the step function equation. Therefore, the linear plateau model was employed to determine the pattern of subterranean turion production in increasing SD.

The plants began forming rhizomes after 20 SD (data not shown) but subterranean turions were not produced until hydrilla was exposed to 24 SD. The linear plateau regression model calculated that after 20 days in SD conditions, hydrilla increased subterranean turion production at the rate 0.17 subterranean turions per day (Figure 2). There is no  $R^2$  value to determine the fit of the equation in non-linear regression, but the slope of the linear plateau model had the lower and upper limits of the confidence interval (CI) above zero [ $0.12 \leq b \leq 0.22$ ] indicating that the slope is significantly greater than zero at the 95% level.

## STUDY 2 (FERTILIZED)

Analysis of variance indicated there was a significant ( $p < 0.05$ ) difference in shoot biomass, subterranean turion number, and subterranean turion biomass among treatments (Table 1).

The first 40 means were grouped as non-inductive and contrasted with the inductive means ( $SD > 40$ ). The shoot biomass of the second part of the study was significantly higher ( $p < 0.05$ ) and the estimated difference between the two groups was calculated (Table 1).

Regression was carried out to predict the change in shoot biomass due to the change in SD conditions (Figure 1). In contrast to the first year study, the shoot biomass of hydrilla increased with increasing SD in the second study which is most likely due to the addition of fertilizer in the second study.

After 20 SD days there was a decrease in the shoot biomass of unfertilized plants (study 1) which coincided with the onset of subterranean turion production. The fertilized plants, on the other hand, began increasing shoot biomass. Since we assume there were ample nutrients in study 2, the vegetative growth may have been prolonged due to less nutrient stress. Due to the delay in reproduction, plants may have had a longer period to increase top growth, causing an increase in shoot biomass.

Analysis of variance indicated that subterranean turion production for the second study was significantly influenced by the duration of SD ( $p < 0.05$ ). Orthogonal contrasts were carried out to determine how the inductive means differed from non-inductive treatment means in their effect on subterranean turion number and weight. The number and biomass of the subterranean turions were significantly greater in the second part of the study than in the first part. The estimated differences between the two treatment groups are listed (Table 1). For further analysis, non-linear regression was carried out to predict the change in subterranean turion production with increasing SD. The linear plateau model calculated that 38 days of exposure to SD conditions were critical for subterranean turion induction in the fertilized plants. After 38 SDs, hydrilla increased subterranean turion production at the rate of 0.89 subterranean turions per day (Figure 2). The slope was significant and positive for the linear plateau model with a CI of [ $0.63 \leq b \leq 1.16$ ].

A possible mechanism that may explain the difference in shoot biomass and subterranean turion production in Study 2 is possible differences in endogenous hormonal regulation. Several researchers have demonstrated that interaction among phytohormones are involved in asexual reproduction in plants (Mohr and Schopfer 1995). Turion formation in *Spirodela polyrrhiza* was found to be controlled by ABA and cytokinins (Smart and Trewas 1983a, Smart and Trewas 1983b). Although ABA is not directly responsible for subterranean turion production in hydrilla, it seems to play an indirect role (MacDonald, 1994). Rhizome formation in dioecious hydrilla depends on a the SD stimulus, and ABA may be produced after rhizome differentiation to maintain dormancy and induce swelling of the rhizome tip (MacDonald 1994).

The experiments reported here showed that 20 to 24 SD stimulate subterranean turion production in unfertilized




Figure 2. The influence of SD exposure period on the subterranean turion production (number per container) in unfertilized and fertilized hydrilla.

plants, and fertilized plants will begin producing subterranean turions after 38 to 44 days of SD conditions. The fertilized plants were more vigorous throughout the study and increased in shoot biomass throughout the study in contrast to the unfertilized plants. The number of subterranean turions produced and the rate of production were significantly higher in fertilized plants, but the onset of turion production was delayed. We also found that once subterranean turions were induced there was no statistically significant difference in turion biomass between the fertilized and non-fertilized plants (data not shown).

The results from these experiments may help in planning more effective and economically feasible herbicide treatment strategies which could be accomplished by applying herbicides or other management strategies at a time that would prevent subterranean turion formation.

### ACKNOWLEDGEMENTS

This research was supported by the Center for Aquatic Plants, the Agronomy Department at the University of Florida, and the USDA/ARS - IFAS/University of Florida Cooperative Agreement.

### LITERATURE CITED

Haller, W. T. 1976. Hydrilla: A New and Rapidly Spreading Aquatic Weed Problem. Circular S-245. Florida Cooperative Extension Service, IFAS, Gainesville, FL 13 pp.

- Klaine, S. J. and C. H. Ward. 1984. Environmental and chemical control of vegetative dormant bud production in *Hydrilla verticillata* Royle. *Ann. of Bot.* 53(4): 503-14.
- Langeland, K. A. 1990. Hydrilla, A Continuing Problem in Florida Waters. Circular No. 884. Cooperative Extension Service, IFAS, Gainesville, FL 21 pp.
- MacDonald, G. E. 1994. Physiological Aspects of Hydrilla [*Hydrilla verticillata* (L.f.) Royle] Growth and Reproduction. Ph.D. Dissertation. University of Florida, Gainesville, FL 77 pp.
- McFarland, D. G. and J. W. Barko. 1990. Temperature and day length effects on growth and tuber formation in Hydrilla. *J. Aquat. Plant Manage.* 28: 15-19.
- Miller, J. D., W. T. Haller, and M. S. Glenn. 1993. Turion production by dioecious hydrilla in North Florida. *J. Aquat. Plant Manage.* 31: 101-105.
- Mitra, E. 1955. Contributions to our knowledge of Indian freshwater plants: I. On some aspects of the structure and life history of *Hydrilla verticillata* with notes on its autecology. *J. of the Asiatic Society.* 21(1): 1-16.
- Mohr, H. and P. Schopfer. 1995. *Plant Physiology*. Springer-Verlag, Berlin. pp. 409-421.
- Netherland, M. D. 1997. Turion ecology of hydrilla. *J. Aquat. Plant Manage.* 35: 1-10.
- Smart, C. M. and A. J. Trewavas. 1983a. Abscisic-acid-induced turion formation in *Spirodela polyrrhiza* L. I. Production and development of the turion. *Plant Cell Environ.* 6: 507-514.
- Smart, C. M. and A. J. Trewavas. 1983b. Abscisic-acid-induced turion formation in *Spirodela polyrrhiza* L. II. Ultrastructure of the turion; a stereo logical analysis. *Plant Cell Environ.* 6: 515-522.
- Spencer, D. F. and L. W. J. Anderson. 1986. Photoperiod responses in monoecious and dioecious *Hydrilla verticillata*. *Weed Sci.* 34: 551-557.
- Van, T. K., W. T. Haller, and G. Bowes. 1978. Some aspects of the competitive biology of hydrilla. *Proc. Eur. Weed Res. Soc.* 5: 1-8.
- Van, T. K. and K. K. Steward. 1990. Longevity of monoecious hydrilla propagules. *J. Aquat. Plant Manage.* 28: 74-76.
- Yeo, R. R., R. H. Falk, and J. R. Thurston. 1984. The morphology of hydrilla [*Hydrilla verticillata* (L.f.) Royle]. *J. Aquat. Plant Manage.* 22: 1-17.