

# Erratum

In the last issue of *Journal of Aquatic Plant Management*, 52#1, January 2014, in the article “Response of invasive floating plants and nontarget emergent plants to foliar applications of imazamox and penoxsulam by Christopher

R. Mudge and M. D. Netherland, page 4, the following table appeared incorrectly. Below is the corrected version. The revised table will be added to the original article when this manuscript is published online.

Table 2. DAYS TO 10% OR GREATER VISUAL INJURY OF FLOATING TARGET AND EMERGENT NON-TARGET AQUATIC PLANT SPECIES TREATED WITH FOLIAR HERBICIDE AND HERBICIDE COMBINATION TREATMENTS.

| Treatment              | Floating target plants |              | Emergent nontarget plants |                   |                     |                    |                  |
|------------------------|------------------------|--------------|---------------------------|-------------------|---------------------|--------------------|------------------|
|                        | Waterhyacinth          | Waterlettuce | Lanceleaf arrowhead       | Jointed spikerush | Gulfcoast spikerush | California bulrush | Softstem bulrush |
| I + S <sup>a,b</sup>   | 10 <sup>c</sup>        | 7            | 14                        | —                 | —                   | —                  | —                |
| I + C + S              | 4                      | 3            | 7                         | 14                | —                   | —                  | —                |
| I + E <sup>d</sup> + S | N/A                    | N/A          | 7                         | 14                | 14                  | —                  | 14               |
| I + F + S              | 4                      | 2            | 7                         | —                 | —                   | 7                  | 7                |
| I + 2 S                | 7                      | 7            | N/A                       | N/A               | N/A                 | N/A                | N/A              |
| P + S                  | 10                     | 7            | 7                         | —                 | —                   | —                  | —                |
| P + C + S              | 4                      | 3            | 7                         | 14                | —                   | 14                 | 14               |
| P + E <sup>d</sup> + S | N/A                    | N/A          | 7                         | 14                | 14                  | 14                 | 14               |
| P + F + S              | 4                      | 2            | 7                         | 14                | —                   | 7                  | 7                |
| P + 2 S                | 7                      | 7            | N/A                       | N/A               | N/A                 | N/A                | N/A              |
| 2,4-D + S              | N/A                    | N/A          | 3                         | 3                 | 3                   | 3                  | 3                |
| 2,4-D + D + S          | N/A                    | N/A          | 1                         | 1                 | 1                   | 1                  | 1                |

<sup>a</sup> Abbreviations: I, imazamox; P, penoxsulam; C, carfentrazone; E, endothall; F, flumioxazin; S, surfactant; D, diquat.

<sup>b</sup> Herbicide (ga.i.ha<sup>-1</sup>) and surfactant application rates: I, 70.1; P, 35.0; C, 16.6; E, 37.1; F, 17.9; S, 1.0% v/v; 2 Surf, 1% plus 0.5% v/v; 2,4-D, 2130.6 (alone) and 1065.3 (combination with D); D,560.7.

<sup>c</sup> Injury indicates number of days until injury was ≥10%; N/A indicates that the herbicide was not applied to the given plant species; — indicates that plant injury never exceeded 10%.

<sup>d</sup> Endothall and 2,4-D were applied as g a.e. ha<sup>-1</sup>.