

Ecological Impact of Grass Carp: A Review of the Available Data

ERIC D. DIBBLE AND KATYA KOVALENKO

ABSTRACT

The exotic grass carp (*Ctenopharyngodon idella*) has been used for almost a half a century in the United States as a biological agent to control and manage aquatic plants. This long-lived generalist herbivore consumes large amounts of vegetation and can considerably alter habitat and impact aquatic communities. We conducted a literature review to determine whether previous studies adequately addressed ecological impacts of grass carp and their underlying mechanisms. Our goal was to identify strengths and limitations of ecological assessment in the literature and suggest a trajectory of future research. The review yielded 1,924 citations on grass carp; however, data on ecological interactions were limited, and most research emphasized the biology of grass carp or eradication of aquatic plants rather than ecological mechanisms responsible for ecosystem-wide impacts. Very few studies addressed effects on habitat complexity or community-structuring processes. We provide a comprehensive tabulated overview of feeding preferences and environmental impacts of grass carp. We argue that ecology is paramount to evaluating grass carp impacts and thorough understanding of these impacts is essential for the appropriate management of aquatic communities. Current knowledge is not sufficient to accurately predict long-term effects of grass carp on freshwater ecosystems. We advise a more cautious approach to developing guidelines for grass carp use.

Key words: aquatic plant management, *Ctenopharyngodon idella*, habitat alteration, literature review.

INTRODUCTION

During the last 45 years, a considerable amount of information has been published on grass carp, *Ctenopharyngodon idella* (Val.). Reports of grass carp use in management started to proliferate in the mid 1970s, approximately 10 years after the first grass carp were stocked into Arkansas reservoirs with the hope that this large exotic herbivore would control growth of noxious aquatic plants (Pierce 1983). These first reports emphasized the biology and physiology of the fish and its feasibility as a tool for weed control (Sills 1970, Stott and Robson 1970, Shireman and Maceina 1981, Shireman 1984, Wiley et al. 1984, 1987, Leslie et al. 1987). Much of this literature was stimulated by the apparent success and efficiency grass carp demonstrated in reducing and even eradi-

cating entire macrophyte communities. Encouraged by the enthusiasm of finding an effective management tool for an ever-increasing problem of invasive plant infestations, this emphasis prevailed in the literature for more than a decade. By the mid-to-late 1980s, the interest shifted toward addressing questions of the direct impacts these herbivores had on environmental parameters other than targeted aquatic plants.

Concurrently, literature was developing in the discipline of aquatic ecology that addressed indirect effects on the structuring of aquatic communities. Much of this literature emphasized effects of exotic introductions and dominant organisms on community structure (Pimm 1987, Ross 1991, Simberloff 1990, Soule 1990, Vitousek 1990) and provided evidence that a keystone species can alter environmental conditions and mediate community processes (Paine 1966), along with other indirect mechanisms responsible for altering community structure (Kerfoot and Sih 1987). Many of these mechanisms involve multi-species interactions, such as competition, predator-prey interactions, and aquatic plants as ecologically important components within aquatic ecosystems (Jeppesen et al. 1997). Submerged macrophytes are important for water quality, nutrient dynamics, and invertebrate-fish interactions (Jeppesen et al. 1997 and references therein). Spatial heterogeneity across a landscape is important for ecological interactions (Palmer and Poff 1997, Schaffer 1981). Like a landscape, a waterscape exhibits similar heterogeneity due to the differences in spatial complexity and structural habitat provided by aquatic plants. Habitat heterogeneity in aquatic systems mediates mechanisms responsible for determining trophic interactions, animal distribution (Werner and Hall 1979, Savino and Stein 1982, Mittelbach 1988, Diehl 1993), and community composition, such as species diversity and abundance (Gillinsky 1984, Adams and DeAngelis 1987). It was previously thought that macrophytes provide few functions other than physical structure, but now it is clear they are grazed by a variety of animals (Lodge et al. 1997) and also indirectly contribute to lake primary productivity by supporting large biomass of periphyton (Wetzel 2001). Vegetated habitats provide abundant food reserves for mammals, waterfowl, amphibians, reptiles, fish, and invertebrates (Swanson and Meyer 1973, Pardue and Nielsen 1979, Gillinsky 1984, Keast 1984, Eldridge 1990, Fredrickson and Laubhan 1996). Grass carp can cause major reduction or complete elimination of aquatic plants when introduced into aquatic systems (Van Dyke et al. 1984, Wiley et al. 1987, Klussman et al. 1988). Diversity of aquatic plant communities can be reduced considerably, even at carp

Department of Wildlife and Fisheries, P.O. Box 9690, Mississippi State Univ., Mississippi State, MS 39762. Received for publication September 2, 2007 and in revised form September 9, 2008.

densities typically required to control a target plant species (Stott and Robson 1970, Stott 1977, Fowler and Robson 1978, Shireman and Maceina 1981). Because such changes in a plant community significantly alter the habitat and interactions of many aquatic organisms, it is prudent to first develop a thorough understanding of how an exotic herbivorous fish can influence the aquatic community before assessing its feasibility as a management tool.

Previous reviews have discussed mostly biology and feeding preferences of grass carp, management applications such as stocking rates, and effects on water quality (e.g., Bain 1993, Petr 2000, Cudmore and Mandrak 2004, Pipalova 2006). None of the previous reviews, however, emphasized ecological interactions or attempted to address potential mechanisms responsible for grass carp impacts. The goals of this review are to evaluate the availability of data in previous research on grass carp that attempt to validate community impacts, identify strengths and limitations of this research, and determine potential design and trajectory of future research, and protocol for use of grass carp as a biocontrol agent.

We used mainly BIOSIS previews and Cambridge Scientific Abstracts to search the literature and yielded 1,924 unduplicated citations on grass carp, 166 of which were pertinent to this review. Research topics and the type of response variables quantified in the research were summarized, and relative prevalence of each topic was determined by calculating the frequency of its occurrence. Publications were evaluated on the basis of whether they attempted to: (1) quantify impacts on community structure and process; (2) measure response variables at habitat, population and/or individual levels; and (3) investigate direct and indirect impacts on the aquatic community. A direct impact was defined as a single-level interaction such as immediate alteration of plant abundance due to grass carp feeding activities; indirect impacts were defined as multi-level interactions, such as changes in zooplankton composition as a result of the increase in phytoplankton abundance due to accelerated nutrient loading facilitated by grass carp foraging activities (Miller and Kerfoot 1987).

RESULTS AND DISCUSSION

Our review of the literature suggested that data are limited for a thorough assessment of grass carp impacts on the aquatic community. We found that most research is related to the eradication of aquatic plants rather than ecological mechanisms potentially responsible for the impacts. Limited data are available to validate long-term consequences and indirect impacts of system-wide introductions of grass carp. Fewer than 50 citations addressing grass carp impacts referred to the term “ecology,” and of these, only a few (<10) inferred ecological processes, yet did not validate causal mechanisms (Table 1). The two most common response variables measured in studies investigating potential impacts by the carp on aquatic communities were direct impact on aquatic plants and indirect alteration of water quality. Less than 2% of all studies investigated structural alteration of habitat or the dynamic processes responsible for structuring the aquatic community.

Direct impacts

Aquatic plant abundance

Changes in plant abundance and community composition occur due to immediate foraging activities as well as alteration of water transparency, disturbance of the sediment, and deposition of fecal matter by grass carp. Grass carp introductions may lead to unexpected and undesirable changes in the plant community. Hanlon et al. (2000) reported several cases of hydrilla (*Hydrilla verticillata* [L.f.] Royle) overgrowth after carp introduction. A shift in plant community composition to less palatable/grazing-resistant species has also been observed (Vinogradov and Zolotova 1974, Hanlon et al. 2000, Pipalova 2002). Several studies reported that grass carp eliminated native plants, leaving invasive vegetation intact (Van Dyke 1994, McKnight and Hepp 1995). Limited data are available to quantify the direct impact on non-target plants since many studies focused only on specific plant species targeted for control and did not assess other plants in the community. In addition, the impact on small and less conspicuous plant species was likely unnoticed because studies frequently used aerial photography to estimate changes in plant abundance.

Grass carp can be considered “selective generalists” in their foraging behavior because they eliminate species in order of decreasing palatability (Colle et al. 1978, Van Dyke et al. 1984, Leslie et al. 1987). More than 50 genera of food items, including aquatic macrophytes, algae, invertebrates and vertebrates, were reported to be eaten by grass carp (Table 2). Feeding preference is one of the best-studied aspects of grass carp biology; however, contradictory evidence exists for several plant species this fish consumes (Tables 2 and 3; also see Bonar et al. 1990, Cooke et al. 2005). Hydrilla and pondweeds (*Potamogeton* spp.) were the most common food items reported to be eaten by grass carp, and species most commonly avoided were in the genera *Nymphaea*, *Potamogeton*, *Myriophyllum*, *Nuphar*, and *Typha*. For the same set of macrophyte species, the order of preference changes under different environmental conditions, most likely determined by ease of mastication (Wiley et al. 1986, Pine et al. 1989, Bonar et al. 1990). Cellulose, silica, and iron content were shown to be the best predictors of unpalatability, whereas calcium and lignin were positively correlated with consumption rates (Bonar et al. 1990). Predictions of the order of plant elimination in the field still have a large degree of uncertainty; the situation may be improved by a series of field studies across a range of environmental conditions.

Habitat heterogeneity

Aquatic plants provide habitat heterogeneity important at both system (i.e., among plant beds) and micro (i.e., within plant beds) scales. The level of this heterogeneity is determined by the magnitude of available spatial “patchiness” constituted by unique plant attributes (i.e., stem-leaf morphology and architecture) found in the habitat (Dibble et al. 2006). Variability and juxtaposition among these different habitat attributes may collectively define the ecological value of habitat in a community or ecosystem as a whole (Wiens 1976, Stensen 1980, Holt 1984, Horne and Schneider 1995,

TABLE 1. GRASS CARP IMPACT ON THE COMMUNITY.

Topical emphasis	References ^a
Effects on Community Structure	
Algae biomass/phytoplankton	8, 11, 24, 30, 32, 41, 44, 48, 53, 55, 56, 61, 63, 65, 75, 110, 114, 123, 124, 137, 161
Crayfish	27, 41, 123
Detritus	61, 113
Fish	3, 4, 8, 9, 10, 11, 16, 19, 22, 26, 42, 46, 50, 52, 53, 54, 55, 56, 58, 61, 62, 63, 67, 73, 78, 80, 85, 86, 92, 94, 99, 100, 104, 105, 108, 113, 121, 123, 141, 143, 151, 157
Habitat alteration	3, 10, 11, 16, 43, 50, 112, 114, 119, 139
Grass shrimp	45, 143
Macroinvertebrates	6, 8, 10, 27, 32, 41, 53, 55, 63, 65, 73, 96, 97, 105, 108, 112
Native plants	30
Periphytic algae	30, 32, 115
Phytoplankton diversity	44, 61, 160
Plant abundance	8, 11, 12, 13, 19, 59, 58, 64, 65, 81, 91, 112, 127, 128, 135, 160
Plant biomass	19, 60, 61, 66, 70, 74, 81, 83, 90, 94, 97, 100, 104, 110, 121, 134, 160, 161
Plant composition	104, 127, 150, 161
Plant diversity	11, 12, 13, 19, 48, 81, 91, 97, 100, 161
Plants, percent coverage	55, 58, 60, 63, 64, 97, 108, 112, 135, 137, 155, 158, 160, 161
Standing crop	84
Seasonal effects on plants	19, 71, 72, 76
Snails	6
Zoobenthos	123, 160
Zooplankton	32, 44, 54, 55, 63, 65, 96, 97, 98, 101, 105, 108, 110, 115
Water quality	4, 7, 8, 11, 17, 18, 19, 30, 31, 32, 44, 47, 48, 53, 54, 55, 58, 61, 63, 65, 66, 67, 74, 75, 79, 94, 95, 96, 98, 100, 101, 104, 105, 106, 108, 110, 113, 114, 115, 121, 123, 124, 137, 143, 150, 160
Effects on Community Process	
Carbon flow	8, 19
Eutrophication	96, 114
Food chains	8, 10, 62, 123
Lake productivity	8, 19, 49, 55
Nutrient loading	18, 24, 86, 113, 114, 124
Photosynthesis	8, 53
Primary production	97, 98, 101, 104, 106, 107, 108, 121, 124
Reduction of buffer capacity	18
Shoreline erosion	33
Streamflow	96
Weevill activity	45

^aReference numbers correspond to the number preceding citations in the bibliography listed in the appendix.

Moloney and Levin 1996). Species diversity is positively correlated with habitat complexity, and increasing heterogeneity among habitats at a landscape scale is a common objective in ecosystem management (Bookhout 1996). Direct impacts by grass carp on habitat heterogeneity, even though ecologically important, were not adequately assessed in the research we reviewed.

Similarly, the heterogeneity at a smaller scale within aquatic plants is just as important for habitat structure (Dionne and Folt 1991, Lillie and Budd 1992, Budd et al. 1995), yet grass carp impacts on structure have gone unmeasured. This level of habitat heterogeneity can be determined as spatial complexity (e.g., plant stem density or frequency of interstices; Lynch and Johnson 1989, Dibble et al. 1996) and is equally important to individual organisms (Anderson 1984, Diehl 1988, Dibble and Harrel 1997, Pedlow et al. 2006) and collectively to populations (Persson and Eklov 1995). Spatial complexity in aquatic habitats is important for predator avoidance and ontogenetic niche shifts (Werner et al. 1977, Lodge et al. 1988, Persson and Crowder 1997) and, at least theoretically, greater macrophyte densities allow higher equi-

librium densities of both fish and macroinvertebrates (Diehl and Kornijów 1997). A few investigations on grass carp hypothesized mechanisms of trophic interactions and population dynamics (Bettoli et al. 1991, 1992, 1993), yet no attempt was made to measure heterogeneity at this level or validates the mechanism that explained the effects. Even without complete vegetation removal, preferential feeding may significantly affect habitat heterogeneity.

The problem of reaching the desired habitat heterogeneity is complicated by the inability to accurately predict effective stocking densities. Despite a relatively high number of studies published on this topic, specific stocking densities that would result in complete elimination of a specifically targeted plant cannot accurately be predicted across a range of different plants and environmental conditions. Partial control of vegetation is rarely achieved with grass carp (Petr 2000, Bonar et al. 2002). After studying 38 lakes in Florida, Hanlon et al. (2000) concluded that the relationship between stocking density and plant abundance was unpredictable except for maximum stocking density that consistently resulted in complete eradication of macrophytes. The prob-

TABLE 2. LIST OF FOOD ITEMS (GENERA) REPORTED EATEN BY GRASS CARP.

Genus	Reference ^a
<i>Alisma</i>	132
<i>Alternanthera</i>	112
<i>Anax</i> ^b	113
<i>Azolla</i>	35, 123, 131
<i>Bacopa</i>	142
<i>Bimastur</i> ^b	113
<i>Brasenia</i>	112
<i>Bufo</i> ^b	113
<i>Cabomba</i>	31, 107, 108, 112, 155
<i>Callitriche</i>	15, 123, 132
<i>Carex</i>	123, 126, 142
<i>Ceratophyllum</i>	4, 10, 19, 35, 48, 55, 60, 64, 79, 93, 95, 98, 101, 104, 107, 121, 123, 126, 127, 131, 132, 134, 150, 155
<i>Chara</i>	15, 35, 48, 66, 72, 75, 81, 95, 99, 123, 126, 128, 130, 142, 143, 146, 160, 164
<i>Cladophora</i>	131, 160, 161
<i>Egeria</i>	84, 91, 95, 134
<i>Eichhornia</i>	26, 35, 45, 112, 131
<i>Eleocharis</i>	40, 84, 87, 112, 142, 143, 161
<i>Elodea</i>	15, 19, 71, 81, 93, 95, 99, 104, 107, 121, 123, 126, 131, 143, 163
<i>Ephemera</i> ^b	113
<i>Glyceria</i>	99
<i>Halophila</i>	123
<i>Hydrilla</i>	2, 4, 7, 10, 13, 17, 18, 20, 31, 33, 35, 44, 48, 49, 54, 55, 58, 62, 63, 64, 70, 78, 79, 91, 95, 96, 98, 105, 108, 120, 123, 133, 135, 155, 158
<i>Hydrocharis</i>	123, 126
<i>Hygrophila</i>	35
<i>Juncus</i>	123, 126
<i>Lagarosiphon</i>	119
<i>Lemna</i>	15, 35, 74, 107, 123, 126, 131, 132
<i>Limnobium</i>	131
<i>Limnophila</i>	123
<i>Mayaca</i>	155
<i>Myriophyllum</i>	4, 10, 15, 18, 35, 48, 55, 64, 66, 71, 75, 93, 95, 96, 99, 101, 107, 123, 126, 130, 132, 134, 142, 155
<i>Nuphar</i>	112
<i>Najas</i>	11, 19, 35, 60, 104, 107, 121, 123, 131, 142, 143, 155, 164
<i>Nasturtium</i>	123, 126
<i>Nechamandra</i>	123
<i>Nitella</i>	15, 84, 123
<i>Notropis</i> ^b	113
<i>Nymphaea</i>	131
<i>Ottelia</i>	123
<i>Phragmites</i>	110, 123, 126
<i>Pistia</i>	35, 131
<i>Pithophora</i>	123
<i>Polygonum</i>	99, 142, 143
<i>Potamogeton</i>	11, 12, 15, 18, 19, 54, 66, 71, 72, 74, 75, 83, 84, 93, 95, 96, 99, 104, 107, 121, 123, 126, 127, 128, 130, 131, 132, 143, 150, 161, 164, 166
<i>Ranunculus</i>	99
<i>Rhizoclonium</i>	15
<i>Ruppia</i>	131
<i>Sagittaria</i>	40, 142
<i>Salomia</i>	123
<i>Salvinia</i>	123
<i>Sparganium</i>	123, 132
<i>Spirodela</i>	123
<i>Spirogyra</i>	123, 131
<i>Trapa</i>	101, 123
<i>Typha</i>	99, 123, 126, 132
<i>Utricularia</i>	87, 107, 123
<i>Vallisneria</i>	123
<i>Wolffia</i>	35, 123
<i>Zannichellia</i>	132
<i>Zygnema</i>	160

^aReference numbers correspond to the number preceding citations in the bibliography listed in the appendix.

^bNon-plant food item.

TABLE 3. AQUATIC PLANTS REPORTED AVOIDED BY GRASS CARP.

Plant Genera	Reference ^a
<i>Alternanthera</i>	123
<i>Brasenia</i>	107
<i>Cabomba</i>	131
<i>Ceratophyllum</i>	40, 161, 164
<i>Chara</i>	60
<i>Cladium</i>	96
<i>Echinodorus</i>	107
<i>Egeria</i>	35
<i>Eichhornia</i>	123
<i>Eleocharis</i>	107
<i>Jussiaea</i>	165
<i>Lemna</i>	166
<i>Myriophyllum</i>	54, 81, 123, 131, 161, 164
<i>Najas</i>	40
<i>Nelumbo</i>	107
<i>Nuphar</i>	31, 81, 108, 109
<i>Nymphaea</i>	31, 93, 107, 108, 109, 123
<i>Nymphoides</i>	109, 123
<i>Orontium</i>	107
<i>Phragmites</i>	160
<i>Pistia</i>	123
<i>Polygonum</i>	107, 123
<i>Potamogeton</i>	35, 40, 107, 166
<i>Ranunculus</i>	109, 165
<i>Sagittaria</i>	107, 131
<i>Salvinia</i>	35
<i>Scirpus</i>	107
<i>Typha</i>	31, 107, 108
<i>Vallisneria</i>	48

^aReference numbers correspond to the number preceding citations in the bibliography listed in the appendix.

lem of appropriate vegetation control is exacerbated by the long life span of this fish; grass carp live longer than 10 years, and they have been shown to persist in a lake for at least 2 years after complete removal of the invasive plant it was stocked to control (Kirk et al. 2000, Kirk and Socha 2003). Long persistence in the environment may also hinder future native macrophyte revegetation efforts if the policy or stakeholder interests change.

Early life stages

Most literature discussed only direct impacts relevant to adult grass carp and neglected potentially important direct impact of early life stages of carp on other fishes. Grass carp stocked for plant control is usually >200 mm in length to reduce mortality due to predation (Pierce 1983, Cassani 1996); however, mounting evidence suggests that naturalized populations of grass carp are reproducing in the United States (Connor et al. 1980, Pflieger and Grace 1987, Brown and Coon 1991, Raibley et al. 1995). In fact, one recent study reported juvenile grass carp in two rivers in Oklahoma previously deemed unsuitable for grass carp reproduction (Hargrave and Gido 2004).

Vegetated habitats provide important nursery grounds for many native fishes (Gregory and Powles 1985, Chubb and Liston 1986, Dibble et al. 1996b), and competition for available food and habitat among and between early life stages of

these fishes can be intense and critical to growth and survival (Werner and Hall 1979, Mittelbach 1981, Diehl 1993). Increases in the number of young non-native fish may exacerbate competition and niche overlap. Larval and juvenile grass carp are not herbivorous and have been shown to feed on zooplankton, insect larvae, chironomids, cladocerans, and copepods (Chilton and Muoneke 1992, Cudmore and Mandrak 2004). Young grass carp may therefore alter trophic dynamics within communities by directly competing for food with native fishes and their larvae.

Indirect impacts

Community structure

Secondary effects from a primary disturbance may be stronger in aquatic communities than the more easily measured direct impacts (Kerfoot and Sih 1987). There is indisputable evidence for presence of trophic cascades in aquatic systems (Brett and Goldman 1996), and some of the strongest cascades are observed in benthic habitats of lakes (Shurin et al. 2002). Presence of novel phytophagous fish exceeding gape size of predator has a potential to involve novel energy pathways (e.g., with more energy flowing towards detrital food chain), which may strongly affect rates of nutrient cycling. Increase in nutrient cycling leads to reduced resilience of an ecosystem (Wetzel 2001), often manifested in population explosions and subsequent crashes of dominant taxa. Because macrophytes play an important role in benthic-pelagic coupling by subsidizing pelagic fish populations and affecting nutrient loading (Schindler and Scheuerell 2002), it is natural to expect that effects of grass carp would propagate beyond the littoral zone and change the role of benthic subsidies to the pelagic habitat of lakes. Mitzner (1978) demonstrated that introduction of grass carp resulted in reduced primary production, possibly due to decreased littoral-pelagic coupling (Lodge et al. 1988).

Grass carp may be viewed as a novel keystone species that reduces spatial heterogeneity, and the resulting loss of complex habitat can potentially influence exploitative competition and other interactions among resident species. This is an example of how the indirect effect of a large herbivore may be transmitted through a resource base, consequently altering the community structure. Other behavioral alterations due to the presence of large fish, whether they are direct predators or not, can impact populations (Sih 1980, Sih et al. 1985, Miller and Kerfoot 1987) and potentially alter ecosystem processes. Secondary changes in aquatic vertebrate and invertebrate communities have been documented after grass carp introductions (see next section). Some observed effects are not consistent across different studies, which may be an indication of our limited understanding of the underlying mechanisms.

Macroinvertebrate populations

Aquatic plants form a base for macrophyte-periphyton-grazer complex (Carpenter and Lodge 1986) and support high densities of macroinvertebrates by providing them with food, habitat, and refuge from predation (reviewed in Diehl and Kornijów 1997). Changes in vegetation due to grass carp

foraging lead to decreased macroinvertebrate diversity and abundance (Vinogradov and Zolotova 1974). Due to decrease in attachment substrate typically provided by vegetated habitat, richness and abundance of epiphytic macroinvertebrates can decrease, whereas the number of benthic invertebrate species can increase (Martin and Shireman 1976, Leslie and Kobylinski 1985, Klussman et al. 1988, Kirkagac and Demir 2004). However, Fedorenko and Frazer (1978) reported that benthic invertebrates also decreased after vegetation removal by grass carp, possibly due to direct competition, low food availability, and increased predation due to the loss of refuge.

Fish populations

In general, when vegetated habitats become too spatially complex, fish growth is negatively impacted due to reduced food availability and decreased feeding efficiency (Crowder and Cooper 1982, Savino and Stein 1982, Anderson 1984). Moderate (20%) removal of overly dense vegetation resulted in improved growth of some age classes of bluegill (Olson et al. 1998); however, further reduction in aquatic plant density and the corresponding decrease in spatial complexity results in increased competition for limited food sources and refugia (Mittelbach 1988). Thus, plant levels at either extreme of the density spectrum may decrease fish growth and survival and, ultimately, alter population dynamics within the community (Adams and DeAngelis 1987, Savino et al. 1992, Diehl 1993).

As a general tendency, a decrease in the abundance of fish species are dependent on aquatic plants, such as largemouth bass (*Micropterus salmoides*) and other sunfishes (*Lepomis* spp.), whereas non-phytophyllic species, such as gizzard shad (*Dorosoma cepedianum*) and silversides (*Labidesthes* spp.), tend to increase in abundance (Ware and Gasaway 1978, Klussman et al. 1988, Maceina et al. 1991, Bettoli et al. 1992, Bettoli et al. (1993) observed a radical change in fish community following grass carp introduction, with a decline in the total number of species over a 7-year period (Table 4). Most notable declines were observed for *Lepomis* spp., crappie (*Pomoxis* spp.), brook silversides (*Labidesthes sicculus*), and juvenile largemouth bass. Colle and Shireman (1994) reported that after complete elimination of vegetation by grass carp, fish harvest declined, and several species had disappeared.

In contrast, Killgore et al. (1998) observed no change in the number of littoral species and an increase in the total fish catch after hydrilla control by grass carp, although the mean total length of the largemouth bass declined (Table 4). This study was unusually successful in achieving intermediate levels of vegetation control, a result that may not be easily extrapolated to many other situations.

Other aquatic vertebrates

Previous work has evaluated indirect impacts of grass carp on waterfowl, suggesting that abundance of bird species can be reduced in the presence of grass carp due to shifts in plant community and competition for preferred food plants (Gasaway and Drda 1976, Johnson and Montalbano 1984, 1987, Leslie et al. 1987, McKnight and Hepp 1995). However, no study specifically addressed grass carp impact on other

TABLE 4. IMPACT OF GRASS CARP ON OTHER FISHES.

Topical emphasis	References ^a
Populations	4, 8, 9, 10, 11, 16, 46, 50, 52, 53, 54, 78, 92, 116, 141
Density/abundance	53, 55, 56, 58, 62, 67, 85, 92, 108, 123, 141, 151, 157
Biomass	8, 50, 55, 58, 61, 78, 81, 92, 100
Diversity	50, 55, 56, 62, 100, 157
Production (kg/ha)	53, 80, 123
Total stand. crop (kg/ha)	58, 62, 63, 73, 78, 108, 123
Weight	92, 105
Relative weight	58
Length	92, 105, 157
Distribution	85, 151
Vulnerability to predation	85
Diets	3, 10, 26, 85, 92, 110, 113
Growth	11, 22, 26, 42, 67, 80, 85, 86, 99
Condition	50, 54, 78, 85, 92, 105, 143
Recruitment	42, 67, 78, 92, 99
Reproductive success	62, 73, 92, 123
Male/female ratio	92
Spawning impacts	16, 81, 123
Angler success	19, 50, 62, 94, 99, 104, 121

^aReference numbers correspond to the number preceding citations in the bibliography listed in the appendix.

vertebrates (e.g., aquatic mammals, reptiles, and amphibians). Many of these animals are highly dependent on vegetated habitats for food and protection from predators, and macrophytes are critical to their survival. Salamanders feed on small prey species living in aquatic plants, and larval stages of amphibians rely on these habitats to avoid predation (Brophy 1980, Zaret 1980).

Muskrats (*Ondatra zibethicus*), beaver (*Castor canadensis*), and nutria (*Myocastor coypus*) rely heavily on aquatic plants (Chabreck 1988, Fredrickson and Laubhan 1996), and more research is needed to quantify how changes in aquatic plants alter habitat and distribution of these animals. A better understanding of how grass carp impact animal distribution will improve the management of these ecosystems; this is especially important for controlling exotic species such as nutria.

Indirect effects operate at different temporal and spatial scales. Parameters that validate those mechanisms are not easily measurable and usually require a period of time before response is detected (Miller and Kerfoot 1987), which may explain why these effects are frequently ignored or inadequately tested. Nevertheless, they need to be incorporated into the designs of future experiments attempting to predict and explain how grass carp impact aquatic communities.

Water quality

Aquatic plants decrease sediment resuspension, play an important role in nutrient cycling (Carpenter and Lodge 1986, Barko and James 1997), and create microclimates in the littoral zone (Lodge et al. 1988). Submerged macrophytes are responsible for several positive feedback mechanisms promoting retention of the vegetated state (Scheffer and Jeppesen 1997, Van Donk 1997), particularly important under the conditions of increasing eutrophication. Shallow

lakes provide one of the most dramatic examples of a catastrophic shift when they suddenly change from clear-water vegetated state to an alternative stable state, turbid and un-vegetated, in response to gradual changes in conditions, such as nutrient loading or macrophyte removal (Scheffer et al. 2001).

Detrimental changes in water quality parameters (increase in nitrite, nitrate, phosphate concentrations) following vegetation control by grass carp were reported in most studies that evaluated water quality (Table 5; Shireman and Smith 1983, Kirkagac and Demir 2004). These changes result from sediment resuspension during feeding and fecal matter deposition by carp as well as collapse of mechanisms responsible for maintenance of the vegetated state due to removal of macrophytes. The rate at which aquatic plants are eliminated determines the magnitude of impact (Hestland and Carter 1978, Lembi et al. 1978, Leslie et al. 1983, 1987, Richard et al. 1984, 1985). These changes in water quality are often followed by algal blooms (Vinogradov and Zolotova 1974, Canfield et al. 1983, Klussman et al. 1988, Shireman and Smith 1983), which in most lakes signal a shift to an alternative stable state (Scheffer and Jeppesen 1997). Increased rates of nutrient cycling after resuspension of sediments lead to decreased ecosystem stability (Wetzel 2001). When weighing pros and cons of complete macrophyte eradication, note that these water quality changes are often irreversible on relatively long time scales, even after herbivorous fish are removed (Scheffer et al. 2001).

Other considerations

Many natural freshwater lakes were isolated for some period of time and may harbor genetically distinct and/or locally rare populations of plants and animals. While it is not clear whether other methods of vegetation control or the invasion itself may have an effect just as detrimental as vegetation control by grass carp, the issue of potential impact on such populations needs to be addressed in the future if carp is to be used in systems other than ponds.

According to the invasional meltdown hypothesis, as more non-native species are introduced into the system, populations of resident native species are increasingly disrupted, and the community becomes more susceptible to future invasions (Simberloff and Von Holle 1999, Ricciardi 2001). As the history of biological introductions has shown, these aspects are very difficult to forecast and assess (Simberloff and Stiling 1996). Until proven otherwise, such possibility exists for grass carp introductions; for instance, frequent disturbance of sediment and vegetation may in fact promote further plant invasions. This may be especially true when grass carp is used without simultaneous mitigation of factors causing nuisance species overgrowth (e.g., eutrophication or frequent disturbance).

Current stocking practices

Grass carp are now recorded in 45 states (Mitchell and Kelly 2006). In some states, only triploid carp can be stocked

TABLE 5. PHYSICAL PARAMETERS MEASURED TO DETERMINE IMPACTS BY GRASS CARP ON WATER QUALITY.

Parameter	References ^a
Alkalinity	7, 17, 19, 47, 53, 54, 55, 58, 63, 66, 75, 94, 95, 104, 114, 121, 143
Ammonia nitrogen	55, 75, 108, 113, 114, 123, 137
Calcium	17, 47, 48, 58
Carbonates	11, 44, 55, 58, 63, 66, 75, 95, 115
Chlorophyll	4, 17, 18, 30, 31, 32, 44, 54, 58, 65, 67, 114, 137, 143, 150, 160
CO ₂	110
Conductivity	7, 17, 54, 55, 58, 63, 66, 95, 108, 114, 115
Dissolved oxygen	7, 44, 47, 18, 54, 66, 95, 96, 100, 101, 105, 110, 114, 116
Dissolved organic matter	8, 79, 98, 106, 115
Iron	115, 123
Kjeldahl nitrogen	18, 54, 66, 95, 96, 115
Light compensation point	105
Magnesium	17, 47, 48, 58, 115, 123
Nitrates/Nitrites	19, 31, 44, 47, 48, 54, 55, 65, 75, 104, 113, 114, 137, 160
Oxygen demand	19, 47, 53, 104
pH	7, 17, 44, 47, 48, 53, 54, 63, 66, 95, 96, 104, 105, 108, 113, 114, 143
Phosphate/Phosphorous	11, 17, 18, 19, 31, 44, 47, 48, 54, 55, 58, 61, 63, 65, 67, 75, 79, 98, 100, 104, 106, 108, 113, 114, 115, 123, 124, 137, 160
Potassium	17, 48, 58, 75, 105, 113, 115, 123
Sulfate	47, 55, 105, 115
Sulfur bacteria	48
Tannin-Lignin levels	31, 47, 108
Temperature	54, 113
Total nitrogen	5, 17, 18, 48, 58, 63, 100, 108
Total photosynthetic pigments	53
Turbidity	18, 19, 47, 48, 53, 54, 66, 95, 104, 108, 110, 123, 143
Water clarity/Secchi	55, 58, 98, 114, 137

^aReference numbers correspond to the number preceding citations in the bibliography listed in the appendix.

for vegetation control, and the procedure for ensuring infertility is tightly regulated by the U.S. Fish and Wildlife Service. Only certified triploid grass carp can be possessed west of the Continental Divide and in the Rio Grande Basin (California Laws: Fish and Game Code, Section 6440-6460). Alaska, Maine, Maryland, Massachusetts, Michigan, Minnesota, Montana, Oregon, North Dakota, Rhode Island, Vermont, and Wisconsin banned both diploid and triploid grass carp (reviewed in Mitchell and Kelly 2006), while some states allow possession of diploid grass carp for aquaculture purposes, production of triploids, or even vegetation control.

In states where grass carp possession is regulated, a special permit (from the departments of natural resources) is required for stocking grass carp. Often, however, a free permit can be obtained by anyone who demonstrated a need to control aquatic vegetation. Satisfactory guidelines have been developed from the standpoint of limited knowledge about ecological effects of this introduction (Table 6). These guidelines, if followed closely, may help ensure that, if deleterious effects occur after grass carp introduction, such effects are at least contained to small ponds not connected to any natural aquatic systems (also see Cooke et al. 2005, p. 443). The recommended maximum stocking density is sufficiently low to prevent, in most cases, complete vegetation removal, and adequate precautions are taken with respect to rare and endangered species. Also, this publication provides the pond owner with sufficient information on the importance of aquatic vegetation and possible changes in water quality after vegetation control by grass carp. Due to its generalist habits, grass carp is not well-suited for selective control of invasive aquatic plants when some level of native plants is desired.

Other Asian carps, such as black (*Mylopharyngodon piceus* Richardson), silver (*Hypophthalmichthys molitrix* Valenciennes), and bighead (*H. nobilis* Richardson) have become invasive species of special concern (reviewed in Schofield et al. 2005), and there is a high likelihood that asian carp may become invasive as well. Some populations have already naturalized, and it is now critical to study their effects on

ecosystems as well as make it unlawful to continue stocking diploid grass carp. The importance of changing regulations in states that still allow possession and stocking of diploids cannot be overemphasized (also see Cooke et al. 2005, p. 437).

Recommendation for future research

More research is needed to investigate both direct and indirect impacts of grass carp, with greater emphasis on indirect impacts. Future investigations need to better quantify these impacts within and across vegetated habitats for a better understanding of community processes. Improved measurement of habitat heterogeneity using new descriptors (e.g., complexity indices, fractal dimension) is essential for an accurate assessment of habitat changes and their effect on community. Additional information is also required to determine how grass carp impact multi-species interactions (e.g., interspecific competition and trophic/predator-prey interactions) and individual behavior (e.g., foraging efficiency, predator avoidance, and habitat use). Future investigations need to address impacts of all life stages of grass carp, especially in systems where reproduction of naturalized populations has been confirmed. It is important to study ecosystem effects of grass carp, such as changes in rates of nutrient cycling, nutrient compartmentalization, and habitat coupling.

Data are needed to determine appropriate levels of abundance and diversity at which aquatic plants should be maintained to prevent a shift to turbid state and negative effects on animal communities. This review shows that aquatic macrophytes are important for habitat heterogeneity and ecological stability; however, it is difficult to achieve optimal habitat heterogeneity and/or only target invasive plants with this non-selective biocontrol agent. Further study of how grass carp feeding preferences across a range of environmental conditions influence spatial attributes of aquatic habitats will help predict community responses to changes in habitat heterogeneity. Meanwhile, innovative research is needed to find suc-

TABLE 6. INFORMATION FROM "GRASS CARP IN NEW YORK PONDS", PUBLICATION OF NEW YORK STATE DEPARTMENT OF ENVIRONMENTAL CONSERVATION, DIVISION OF FISH, WILDLIFE & MARINE RESOURCES.

“... approve and issue permits for stocking of up to 15 United States Fish and Wildlife Service certified triploid grass carp [from approved suppliers possessing a valid permit to import and sell triploid grass carp in New York State] per surface acre for aquatic plant management purposes in ponds five (5) acres or less in size which lie wholly within the boundaries of lands privately owned or leased by the individual making or authorizing such treatments if:

- aquatic plants targeted for control significantly impair the intended use(s) of the pond;
- the pond harbors no species of wildlife, fish, shellfish or crustacea identified by the Department as being endangered, threatened or special concern; or any species of plant identified as being endangered, threatened or rare;
- the pond is not contiguous to or part of a New York State regulated freshwater wetland.
- is not an impoundment or natural pond on a permanent stream or a source of a permanent stream.
- at least two years have elapsed since the last stocking of triploid grass carp

“Triploid grass carp are extremely potent plant consumers. If overstocked, they are capable of eradicating all plants from a pond for periods exceeding 10 years. Besides the obvious impact such complete plant removal will have on vegetation-dependent fish and wildlife, total devegetation of a pond can also result in the development of severe algae blooms, foul smells and an overall decline in water clarity. To minimize or prevent such adverse impacts, plant populations should be maintained at approximately 20-30% of the pond’s surface area.

Due to various factors that impact triploid grass carp feeding, it is impossible to precisely predict the exact number of fish to stock to achieve the 20-30% plant coverage target. The only way to prevent excessive plant control is through use of an incremental approach. This approach involves the stocking of triploid grass carp at the stocking rates suggested below, followed by a two-year waiting period for the fish to achieve maximal control. Then, if needed, more fish are added in small increments at two-year intervals until plant populations are reduced to the 20-30% threshold.”

cessful methods to provide more accurate control for different levels of plant growth, abundance, and composition. Detailed evaluation of community responses to differential stocking rates is needed, as are the techniques to control current grass carp populations already released into natural systems.

Understanding ecological impacts is paramount to appropriate management of aquatic communities. Current data are not sufficient to adequately answer important questions about use of grass carp as a biocontrol agent for aquatic plants. The problem of assessing all possible impacts is not unique to grass carp: it has been shown for most introduced biocontrol agents (reviewed in Simberloff and Stiling 1996). The extent to which these approaches are used in future research will determine the amount of knowledge gained relative to understanding the ecological impacts of grass carp on aquatic communities. Until these data are collected, a more conservative approach should be used when developing guidelines for grass carp use.

ACKNOWLEDGMENTS

We appreciate the help provided by J. Cruickshank, E. Forrester, R. Kimbrough, B. LaValley, and J. O'Keefe for their assistance in the retrieval and initial analysis of literature, and S. Harrel and P. Smiley for editorial comments provided on an earlier draft of this manuscript. This manuscript was significantly improved by the suggestions of three anonymous reviewers. The Aquatic Ecosystem Restoration Foundation funded an earlier review of the literature that was incorporated in this manuscript. We thank J. Madsen and the GeoResources Institute at Mississippi State University for support and the U.S.G.S. Invasive Species Science Program for funding this project.

LITERATURE CITED

Adams, S. M. and D. L. DeAngelis. 1987. Indirect effects of early bass-shad interaction on predator population structure and food web dynamics, pp. 103-117. *In*: W. C. Kerfoot and A. Sih (eds). Predation: Direct and Indirect Impacts on Aquatic Communities. University Press of New England, Hanover, NH.

Anderson, O. 1984. Optimal foraging by largemouth bass in structured environments. *Ecology* 65:351-861.

Bain, M. B. 1993. Assessing impacts of introduced aquatic species: grass carp in large systems. *Environ. Manage.* 17:509-516.

Barko, J. W. and W. F. James. 1997. Effects of submerged aquatic macrophytes on nutrient dynamics, sedimentation, and resuspension, pp. 197-214. *In*: E. Jeppesen, M. Søndergaard, M. Søndergaard and K. Christoffersen (eds.). The structuring role of submerged macrophytes in lakes. Springer, NY.

Bettoli, P. W., J. E. Morris and R. L. Noble. 1991. Changes in the abundance of two atherinid species after aquatic vegetation removal. *Trans. Am. Fish. Soc.* 120:90-97.

Bettoli, P. W., M. J. Maceina, R. L. Noble and R. K. Betsill. 1992. Piscivory in largemouth bass as a function of aquatic vegetation abundance. *N. Am. J. Fish. Manage.* 12:509-516.

Bettoli, P. W., M. J. Maceina, R. L. Noble and R. K. Betsill. 1993. Response of a reservoir fish community to aquatic vegetation removal. *N. Am. J. Fish. Manage.* 13:110-124.

Bonar, S. A., H. S. Sehgal, G. B. Pauley and G. L. Thomas. 1990. Relationship between the chemical composition of aquatic macrophytes and their consumption by grass carp, *Ctenopharyngodon idella*. *J. Fish Biol.* 36:149-157.

Bonar, S. A., B. Bolding and M. Divens. 2002. Effects of triploid grass carp on aquatic plants, water quality and public satisfaction in Washington State. *N. Am. J. Fish. Manage.* 21:96-105.

Bookhout, T. A. (ed.). 1996. Research and management techniques for wildlife and habitats. The Wildlife Society, Allen Press, Lawrence, KS. 740 pp.

Brett, M. T. and C. R. Goldman. 1996. A meta-analysis of the freshwater trophic cascade. *P. Natl. Acad. Sci. USA* 93:7723-7726.

Brophy, T. E. 1980. Food habits of sympatric larval *Ambystoma tigrinum* and *Notophthalmus viridescens*. *J. Herpetol.* 14:1-6.

Brown, D. J. and T. G. Coon. 1991. Grass carp larvae in the lower Missouri River and its tributaries. *N. Am. J. Fish. Manage.* 11:62-66.

Budd, J., R. A. Lillie and P. Rasmussen. 1995. Morphological characteristics of the aquatic macrophyte, *Myriophyllum spicatum* L. in Fish Lake, Wisconsin. *J. Freshw. Ecol.* 10(1):19-31.

California Laws: Fish And Game Code. Section 6460. <http://www.aroundthecapitol.com/code/code.html?sec=fgc&code=6440-6460>. April 2, 2009.

Canfield, D. E. Jr., M. J. Maceina and V. J. Shireman. 1983. Effects of Hydrilla and grass carp on water quality in a Florida lake. *Water Resour. Bull.* 19:773-778.

Carpenter, S. R. and D. M. Lodge. 1986. Effects of submersed macrophytes on ecosystem processes. *Aquat. Bot.* 26:341-370.

Cassani, J. R. (ed). 1996. Managing aquatic vegetation with grass carp, a guide for resource managers. American Fisheries Society, Bethesda, MD. 196 pp.

Chabreck, R. H. 1988. Coastal marshes: ecology and wildlife management. Minnesota Press, MN. 139 pp.

Chilton E. W. II and M. I. Muoneke. 1992. Biology and management of grass carp (*Ctenopharyngodon idella*, Cyprinidae) for vegetation control: a North American perspective. *Rev. Fish. Biol. Fish.* 2:283-320.

Chubb, S. L. and C. R. Liston. 1986. Density and distribution of larval fishes in Pentwater Marsh, a coastal wetland on Lake Michigan. *J. Gt. Lakes Res.* 12:332-343.

Colle, D. E., J. V. Shireman and R. W. Rottmann. 1978. Food selection by grass carp fingerlings in a vegetated pond. *Trans. Am. Fish. Soc.* 107:149-152.

Colle, D. E. and J. V. Shireman. 1994. Use of grass carp in two Florida lakes, 1975 to 1994, pp. 111-120. *In*: Proceedings of the Grass Carp Symposium. U.S. Army Corps of Engineers, Vicksburg, MS.

Connor, J. B., R. P. Gallagher and M. F. Chatry. 1980. Larval evidence for natural reproduction of the grass carp *Ctenopharyngodon idella* in the lower Mississippi River. U.S. Fish and Wildlife Service Biological Services Program, FWS/OBS-80/43-14.

Cooke, D. G., E. B. Welch, S. A. Peterson and S. A. Nichols. 2005. Restoration and management of lakes and reservoirs. 3rd edition. CRC Press, Taylor & Francis, Boca Raton, FL. 591 pp.

Crowder, L. B. and W. E. Cooper. 1982. Habitat structural complexity and the interaction between bluegills and their prey. *Ecology* 63:1802-1813.

Cudmore, B. and N. E. Mandrak. 2004. Biological synopsis of grass carp (*Ctenopharyngodon idella*). *Can. Man. Report Fish. Aquat. Sci.* 2705. 44 pp.

Dibble, E. D., G. Dick and K. J. Killgore. 1996a. Measurement of plant architecture in seven aquatic plants. *J. Freshw. Ecol.* 11: 311-318.

Dibble, E. D., K. J. Killgore and S. L. Harrel. 1996b. Assessment of fish-plant interactions, pp. 357-372. *In*: L. E. Miranda and D. R. DeVries (eds.). Multidimensional approaches to reservoir fisheries management. *Am. Fish. Soc.*, Bethesda, MD.

Dibble, E. D. and S. L. Harrel. 1997. Largemouth bass diets in two aquatic plant communities. *J. Aquat. Plant Manage.* 35:74-78.

Dibble, E. D., S. M. Thomaz and A. A. Padial. 2006. Spatial complexity measured at a multi-scale in three aquatic plant species. *J. Freshw. Ecol.* 21:239-247.

Diehl, S. 1988. Foraging efficiency of three fresh-water fishes: effects of structural complexity and light. *Oikos* 53:207-214.

Diehl, S. 1993. Effects of habitat structure on resource availability, diet and growth of benthivorous perch, *Perca fluviatilis*. *Oikos* 67:403-414.

Diehl, S. and R. Kornijów. 1997. Influence of submerged macrophytes on trophic interactions among fish and macroinvertebrates, pp. 24-46. *In*: E. Jeppesen, M. Søndergaard, M. Søndergaard and K. Christoffersen (eds.). The structuring role of submerged macrophytes in lakes. Springer, NY.

Dionne, M. and C. L. Folt. 1991. An experimental analysis of macrophyte growth as fish foraging habitat. *Can. J. Fish. Aquat. Sci.* 48:123-131.

Eldridge, J. 1990. Aquatic invertebrates important for waterfowl production. U.S. Fish and Wildlife Service, Waterfowl Manage. 13:3-7.

Fedorenko, A. Y. and F. J. Fraser. 1978. A review of the biology of grass carp (*Ctenopharyngodon idella*, Val.) and its evaluation as a potential weed control agent in British Columbia. Fisheries and Marine Service Technical Report No. 786. 15p.

Fowler, M. C. and T. O. Robson. 1978. The effects of food preferences and stocking densities of grass carp (*Ctenopharyngodon idella* Val.) on mixed plant communities. *Aquat. Bot.* 5:261-276.

Fredrickson, L. H. and M. K. Laubhan. 1996. Managing wetlands for wildlife, pp. 623-647. *In*: T. A. Bookhout (ed). Research and managements tech-

- niques for wildlife and habitats. The Wildlife Society, Allen Press, Lawrence, KS.
- Gasaway, R. D. and T. F. Drda. 1976. Effects of grass carp introduction on waterfowl habitat. North American Wildlife and Natural Resources Conference 42:73-85.
- Gilinsky, E. 1984. The role of fish predation and spatial heterogeneity in determining benthic community structure. *Ecology* 4:455-468.
- Gregory, R. S. and P. M. Powles. 1985. Chronology, distribution, and sizes of larval fish sampled by light traps in macrophytic Chemung Lake. *Can. J. Zool.* 19:2569-2577.
- Hanlon, S. G., M. V. Hoyer, C. E. Cichra and D. E. Canfield. 2000. Evaluation of macrophyte control in 38 Florida lakes using triploid grass carp. *J. Aquat. Plant Manage.* 38:48-54.
- Hargrave, C. W. and K. B. Gido. 2004. Evidence of reproduction by exotic grass carp in the Red and Washita rivers, Oklahoma. *Southwest. Nat.* 49(1):89-93.
- Hestland, R. S. and C. C. Carter. 1978. Comparative effects of grass carp and selected herbicides on macrophytes and phytoplankton communities. *J. Aquat. Plant Manage.* 16:43-50.
- Holt, R. D. 1984. Spatial heterogeneity, indirect interactions, and the coexistence of prey species. *Am. Nat.* 124:197-229.
- Horne, J. K. and D. C. Schneider. 1995. Spatial variance in ecology. *Oikos* 74:18-26.
- Jeppesen, E., M. Søndergaard, M. Søndergaard and K. Christoffersen (eds.). 1997. The structuring role of submerged macrophytes in lakes. *Ecological Studies*, vol. 131. Springer, NY. 423 pp.
- Johnson, F. A. and F. Montalbano III. 1984. Selection of plant communities by wintering waterfowl on Lake Okeechobee, Florida. *J. Wildl. Manage.* 48:174-178.
- Johnson, F. A. and F. Montalbano III. 1987. Considering waterfowl habitat in hydrilla control policies. *Wildlife Soc. B.* 15:466-469.
- Keast, A. 1984. The introduced aquatic macrophyte, *Myriophyllum spicatum*, as habitat for fish and their invertebrate prey. *Can. J. Zool.* 62:1289-1303.
- Kerfoot, C. W. and A. Sih (eds.). 1987. Predation: direct and indirect impacts on aquatic communities. Univ. Press of New England, Hanover, NH. 366 pp.
- Killgore, K. J., J. P. Kirk and J. W. Foltz. 1998. Response of littoral fishes in Upper Lake Marion, South Carolina following *Hydrilla* control by triploid grass carp. *J. Aquat. Plant Manage.* 36:82-87.
- Kirk, J. P., J. V. Morrow, K. J. Killgore, S. J. De Kozlowski and J. W. Preacher. 2000. Population response of triploid grass carp to declining levels of *Hydrilla* in the Santee Cooper reservoirs, South Carolina. *J. Aquat. Plant Manage.* 38:14-17.
- Kirk, J. P. and R. C. Socha. 2003. Longevity and persistence of triploid grass carp stocked into the Santee Cooper reservoirs of South Carolina. *J. Aquat. Plant Manage.* 41:90-92.
- Kirkagac, M. and N. Demir. 2004. The effects of grass carp on aquatic plants, plankton and benthos in ponds. *J. Aquat. Plant Manage.* 42:32-39.
- Klussman, W. G., R. L. Noble, R. D. Martyn, W. J. Clark, R. K. Betsill, P. W. Bettoli, M. F. Cichra and J. M. Campbell. 1988. Control of aquatic macrophytes by grass carp in Lake Conroe, Texas, and the effect on the reservoir ecosystem. Texas Agriculture Experiment Station MP-1664, College Station, TX.
- Lembi, C. A., B. G. Ritenour, E. M. Iverson and E. C. Fors. 1978. The effects of vegetation removal by grass carp on water chemistry and phytoplankton in Indiana ponds. *Trans. Am. Fish. Soc.* 107:161-171.
- Leslie, A. J. Jr., L. E. Nall and J. M. Van Dyke. 1983. Effects of vegetation control by grass carp on selected water quality variables in four Florida lakes. *Trans. Am. Fish. Soc.* 107:161-171.
- Leslie, A. J. Jr. and G. J. Kobylinski. 1985. Benthic macroinvertebrate response to aquatic vegetation removal by grass carp in a north Florida reservoir. *Fla. Sci.* 48:220-231.
- Leslie, A. J. Jr., J. M. Van Dyke, R. S. Hestland III and B. Z. Thompson. 1987. Management of aquatic plants in multi-use lakes with grass carp (*Ctenopharyngodon idella*), pp. 266-276. *In: G. Redfield* (ed). Lake and reservoir management, vol. III. N. Am. Lake Manage. Soc., Washington, DC.
- Lillie, R. A. and J. Budd. 1992. Habitat architecture of *Myriophyllum spicatum* L. as an index to habitat quality for fish and macroinvertebrates. *J. Freshw. Ecol.* 7:113-125.
- Lodge, D. M., J. W. Barko, D. Strayer, J. M. Melack, G. M. Mittelbach, R. W. Howarth, B. Menge and J. E. Titus. 1988. Spatial heterogeneity and habitat interactions in lake communities, pp. 181-208. *In: S. R. Carpenter* (ed.). Complex interactions in lake communities. Springer-Verlag, NY.
- Lodge D. M., G. Cronin, E. Van Donk and A. J. Froelich. 1997. Impact of herbivory on plant standing crop: comparisons among biomes, between vascular and non-vascular plants, and among freshwater herbivore taxa, pp. 149-174. *In: E. Jeppesen, M. Søndergaard, M. Søndergaard and K. Christoffersen* (eds.). The structuring role of submerged macrophytes in lakes. Springer, NY.
- Lynch, W. E. Jr. and D. L. Johnson. 1989. Influences of interstice size, shade, and predators on the use of artificial structures by bluegills. *N. Am. J. Fish. Manage.* 9:219-225.
- Maccina, M. J., P. W. Bettoli, W. G. Klussmann, R. K. Betsill and R. L. Noble. 1991. Effect of aquatic macrophyte removal on recruitment and growth of black crappies and white crappies in Lake Conroe Texas. *N. Am. J. Fish. Manage.* 11:556-563.
- Martin, R. G. and J. V. Shireman. 1976. A quantitative sampling method for hydrilla-inhabiting macroinvertebrates. *J. Aquat. Plant Manage.* 14:16-19.
- McKnight, S. K. and G. R. Hepp. 1995. Potential effect of grass carp herbivory on waterfowl foods. *J. Wildl. Manage.* 59:720-727.
- Miller, T. E. and W. C. Kerfoot. 1987. Redefining indirect effects, pp. 33-37. *In: C. W. Kerfoot and A. Sih* (eds). Predation: direct and indirect impacts on aquatic communities. Univ. Press of New England.
- Miranda, L. E., W. L. Shelton and T. D. Bryce. 1984. Effects of water level manipulation on abundance, mortality and growth of young-of-year largemouth bass in West Point Reservoir, Alabama-Georgia. *N. Am. J. Fish. Manage.* 4:314-320.
- Mitchell, A. J. and A. M. Kelly. 2006. The public sector role in the establishment of grass carp in the United States. *Fisheries* 31:113-121.
- Mittelbach, G. G. 1981. Foraging efficiency and body size: a study of optimal diet and habitat use by bluegills. *Ecology* 62:1370-1386.
- Mittelbach, G. G. 1988. Competition among refuging sunfishes and effects of fish density on littoral zone invertebrates. *Ecology* 69:614-623.
- Mitzner, L. 1978. Evaluation of biological control of nuisance aquatic vegetation by grass carp. *Trans. Am. Fish. Soc.* 107(1):135-145.
- Moloney, K. A. and S. A. Levin. 1996. The effects of disturbance architecture on landscape-level population dynamics. *Ecology* 77:375-394.
- Olson, M. H., S. R. Carpenter, P. Cunningham, S. Gafny, B. R. Herwig, N. P. Nibbelink, T. Pellett, C. Storlie, A. S. Trebitz and K. A. Wilson. 1998. Managing macrophytes to improve fish growth: a multi-lake experiment. *Fisheries* 23:6-12.
- Paine, R. T. 1966. Food web complexity and species diversity. *Am. Nat.* 100:65-75.
- Palmer, M. A. and N. L. Poff. 1997. The influence of environmental heterogeneity on patterns and processes in streams. *J. N. Am. Benthol. Soc.* 16:169-173.
- Pardue, G. B. and L. A. Nielsen. 1979. Invertebrate biomass and fish production in ponds with added attachment surface, pp. 34-43. *In: D. L. Johnson and R. A. Stein* (eds.). Response of fish to habitat structure in standing water. *Am. Fish. Soc., North Central Division, Special Publication* 6. Bethesda, MD.
- Pedlow, C. L., E. D. Dibble and K. D. Getsinger. 2006. Littoral habitat heterogeneity and shifts in plant composition relative to a fall whole-lake fluridone application in Perch Lake, MI. *J. Aquat. Plant Manage.* 44:26-31.
- Persson, L. and P. Eklov. 1995. Prey refuges affecting interactions between piscivorous perch and juvenile perch and roach. *Ecology* 76:70-81.
- Persson, L. and L. B. Crowder. 1997. Fish-habitat interactions mediated via ontogenetic niche shifts, pp. 3-23. *In: E. Jeppesen, M. Søndergaard, M. Søndergaard and K. Christoffersen* (eds.). The structuring role of submerged macrophytes in lakes. Springer, NY.
- Petr, T. 2000. Interaction between fish and aquatic macrophytes in inland waters, pp. 16-27. *In: FAO Fisheries Technical Paper* 396. Rome, FAO. 185 pp.
- Pflieger, W. L. and T. B. Grace. 1987. Changes in the fish fauna of the lower Missouri River, 1940-1983, pp. 166-177. *In: W. J. Matthews and D. C. Heins* (eds.). Community and evolutionary ecology of North American stream fishes. Univ. of Oklahoma Press, Norman, OK.
- Pierce, B. A. 1983. Grass carp status in the United States: a review. *Environ. Manage.* 7(2):151-160.
- Pimm, S. L. 1987. Determining the effects of introduced species. *Trends Ecol. Evol.* 2:106-108.
- Pine, R. T., L. W. J. Anderson and S. S. O. Hung. 1991. Effects of static versus flowing water on aquatic plant preferences of triploid grass carp. *Trans. Am. Fish. Soc.* 118:336-344.
- Pipalova, I. 2006. A review of grass carp use for aquatic weed control and its impact on water bodies. *J. Aquat. Plant Manage.* 44:1-12.
- Raibley, P. T., D. Blodgett and R. E. Sparks. 1995. Evidence of grass carp (*Ctenopharyngodon idella*) reproduction in the Illinois and Upper Mississippi Rivers. *J. Freshw. Ecol.* 10:65-74.
- Ricciardi, A. 2001. Facilitative interactions among aquatic invaders: is an "invasional meltdown" occurring in the Great Lakes? *Can. J. Fish. Aquat. Sci.* 58:2513-2525.

- Richard, D. I., J. W. Small Jr. and J. A. Osborne. 1984. Phytoplankton responses to reduction and elimination of submerged vegetation by herbicides and grass carp in four Florida lakes. *Aquat. Bot.* 20:307-319.
- Richard, D. I., J. W. Small and J. A. Osborne. 1985. Response of zooplankton to the reduction and elimination of submerged vegetation by grass carp and herbicide in four Florida lakes. *Hydrobiologia* 123:97-108.
- Ross, S. T. 1991. Mechanisms structuring fish assemblages: are there lessons from introduced species? *Environ. Biol. Fish.* 30:359-368.
- Savino, J. F. and R. A. Stein. 1982. Predator-prey interaction between largemouth bass and bluegills as influenced by simulated, submersed vegetation. *Trans. Am. Fish. Soc.* 11:255-266.
- Savino, J. F., E. A. Marschall and R. A. Stein. 1992. Bluegill growth as modified by plant-density: an exploration of underlying mechanisms. *Oecologia* 89:153-160.
- Schaffer, W. M. 1981. Ecological abstractions: the consequences of reduced dimensionality in ecological models. *Ecol. Monogr.* 51:383-401.
- Scheffer, M. and E. Jeppesen. 1997. Alternative stable states, pp. 397-406. *In: E. Jeppesen, M. Søndergaard, M. Søndergaard and K. Christoffersen (eds.). The structuring role of submerged macrophytes in lakes.* Springer, NY.,
- Scheffer, M., S. Carpenter, J. A. Foley, C. Folke and B. Walker. 2001. Catastrophic shifts in ecosystems. *Nature* 413:591-596.
- Schindler, D. E. and M. D. Scheuerell. 2002. Habitat coupling in lake ecosystems. *Oikos* 98:177-189.
- Schofield, P. J., Williams, J. D., Nico, L. G., Fuller, P. and M. R. Thomas. 2005. Foreign Nonindigenous Carps and Minnows (Cyprinidae) in the United States—A guide to their Identification, Distribution, and Biology: U.S. Geological Survey Scientific Investigations Report 2005-5041. 103 pp.
- Shireman, J. V. 1984. Control of aquatic weeds with exotic fishes, pp. 302-312. *In: W. R. Courtenay Jr. and J. R. Stauffer Jr. (eds.). Distribution, biology and management of exotic fishes.* The John Hopkins Univ. Press. Baltimore, MD.
- Shireman, J. V. and M. J. Maceina. 1981. The utilization of grass carp, *Ctenopharyngodon idella* Val., for hydrilla control in Lake Baldwin, Florida. *J. Fish Biol.* 19:629-636.
- Shireman, J. V. and C. R. Smith. 1983. Synopsis of the biological data on the grass carp, *Ctenopharyngodon idella* (Cuvier and Valenciennes, 1944). Food and Agriculture Organization of the United Nations, FAO Fisheries Synopsis Number 135, Rome.
- Shurin, J. B., E. T. Borer, E. W. Seabloom, K. Anderson, C. A. Blanchette, B. Broitman, S. D. Cooper and B. S. Halpern. 2002. A cross-ecosystem comparison of the strength of trophic cascades. *Ecol. Lett.* 5:785-791.
- Sih, A. 1980. Optimal behavior: can foragers balance two conflicting demands? *Science* 210:1041-1043.
- Sih, A. P. Crowley, M. Mcpeck, J. Petrank and K. Strohmeier. 1985. Predation, competition, and prey communities: a review of field experiments. *Ann. Rev. Ecol. Syst.* 16:269-305.
- Sills, J. B. 1970. A review of herbivorous fish for weed control. *Prog. Fish Cult.* 32(3):158-161.
- Simberloff, D. 1990. Community effects of biological introductions and their implication for restoration, pp. 128-136. *In: D. R. Towns, C. H. Daugherty and I. A. Atkinson (eds.). Engineered organisms in the environment: scientific issues.* Am. Soc. Microbiol. Washington, DC.
- Simberloff, D. and P. Stiling. 1996. How risky is biological control? *Ecology* 77(7):1965-1974.
- Simberloff, D. and B. Von Holle. 1999. Positive interactions of nonindigenous species: invasional meltdown? *Biol. Invasions* 1:21-32.
- Soule, M. E. 1990. The onslaught of alien species, and other challenges in the coming decades. *Conserv. Biol.* 4:233-239.
- Stensen, N. C. 1980. Spatial heterogeneity and population stability: some evolutionary consequences. *Oikos* 35:165-184.
- Stott, B. 1977. On the question of the introduction of the grass carp (*Ctenopharyngodon idella* Val.) into the United Kingdom. *Fish. Manage.* 8:63-71.
- Stott, B. and T. O. Robson. 1970. Efficiency of grass carp (*Ctenopharyngodon idella* Val.) in controlling submerged water weeds. *Nature* 226:870.
- Swanson, G. A. and M. I. Meyer. 1973. The role of invertebrates in the feeding ecology of Anatinae during the breeding season, pp. 143-165. *In: Proceedings of the Waterfowl Habitat Symposium.* Moncton, New Brunswick, Canada.
- Van Donk, E. 1997. Switches between clear and turbid water states in a biomanipulated lake (1986-1996): the role of herbivory on macrophytes, pp. 290-297. *In: E. Jeppesen, M. Søndergaard, M. Søndergaard and K. Christoffersen (eds.). The structuring role of submerged macrophytes in lakes.* Springer, NY.
- Van Dyke, J. M., A. J. Leslie Jr. and L. E. Nall. 1984. The effects of the grass carp on aquatic macrophytes of four Florida lakes. *J. Aquat. Plant Manage.* 22:87-95.
- Van Dyke, J. M. 1994. Long-term use of grass carp for aquatic plant control in Deer Point lake, Bay County, Florida, pp. 146-150. *In: Proceedings of the Grass Carp Symposium.* U.S. Army Corps of Engineers, Vicksburg, MS.
- Vinogradov, V. K. and Z. K. Zolotova. 1974. The influence of the grass carp on aquatic ecosystems. *Hydrobiol. J.* 10(2):72-78.
- Vitousek, P. M. 1990. Biological invasions and ecosystem processes: toward an integration of population biology and ecosystem studies. *Oikos* 57:7-13.
- Ware, F. J. and R. D. Gasaway. 1978. Effects of grass carp on native fish population in two Florida lakes. *Proceedings of the Annual Conference of the Southeastern Association of Fish and Wildlife Agencies* 30:324-335.
- Werner, E. E., D. J. Hall, D. R. Laughlin, D. J. Wagner, L. A. Wilmann and F. C. Funk. 1977. Habitat partitioning in a freshwater fish community. *J. Fish. Res. Board Can.* 34:360-370.
- Werner, E. E. and D. J. Hall. 1979. Foraging efficiency and habitat switching in competing sunfishes. *Ecology* 60:256-264.
- Wetzel, R. G. 2001. *Limnology: lake and river ecosystems.* 3rd edition. Academic Press, San Diego, CA. 1006 pp.
- Wiens, J. A. 1976. Population responses to patchy environments. *Ann. Rev. Ecol. Syst.* 7:81-120.
- Wiley, M. J., R. W. Gorden, S. W. Waite and T. Powless. 1984. The relationship between aquatic macrophytes and sport fish production in Illinois pond: a simple model. *N. Am. J. Fish. Manage.* 4:111-119.
- Wiley, M. J., S. M. Pescitelli and L. D. Wike. 1986. The relationship between feeding preferences and consumption rates in grass carp and grass carp x bighead carp hybrids. *J. Fish Biol.* 29:507-514.
- Wiley, M. J., P. P. Tazik and S. T. Sobaski. 1987. Controlling aquatic vegetation with triploid grass carp. Circular 57. Illinois Natural History Survey, Champaign, IL.
- Zaret, T. M. 1980. *Predation and freshwater communities.* Yale Univ. Press. New Haven, CT. 187 pp.

APPENDIX

Bibliography (Grass Carp Papers Reviewed)

- 1) Pierce, B. A. 1983. Grass carp status in the United States: a review. *Environ. Manage.* 7(2):151-160.
- 2) Richard, D. I., J. W. Small and J. A. Osborne. 1985. Response of zooplankton to the reduction and elimination of submerged vegetation by grass carp and herbicide in four Florida lakes. *Hydrobiologia* 123:97-108.
- 3) Petridis, D. 1990. The influence of grass carp on habitat structure and its subsequent effect on the diet of tench. *J. Fish Biol.* 36:533-544.
- 4) Bettoli, P. W., T. Springer and R. L. Noble. 1990. A deterministic model of the response of threadfin shad to aquatic macrophyte control. *J. Freshw. Ecol.* 5(4):445-454.
- 5) Allen, S. K. Jr. and R. J. Wattendorf. 1987. Triploid grass carp: status and management implications. *Fisheries* 12(4):20-24.
- 6) Van Schayck, C. P. 1986. The effect of several methods of aquatic plant control on two *Bilharzia*-bearing snail species. *Aquat. Bot.* 24:303-309.
- 7) Small, J. W. Jr, D. I. Richard and J. A. Osborne. 1985. The effects of vegetation removal by grass carp and herbicides on the water chemistry of four Florida lakes. *Freshw. Biol.* 15:587-596.
- 8) Ewel, K. C. and T. D. Fontaine III. 1982. Effects of white amur (*Ctenopharyngodon idella*) on a Florida lake: a model. *Ecol. Model.* 16:251-273.
- 9) Swar, D. B. and T. B. Gurung. 1988. Introduction and cage culture of exotic carps and their impact on fish harvested in Lake Begnas, Nepal. *Hydrobiologia* 166:277-283.
- 10) Bettoli, P. W., J. E. Morris and R. L. Noble. 1991. Changes in the abundance of two atherinid species after aquatic vegetation removal. *Trans. Am. Fish. Soc.* 120:90-97.
- 11) Buck, D. H., R. J. Baur and C. R. Rose. 1975. Comparison of the effects of grass carp and the herbicide diuron in densely vegetated pools containing golden shiners and bluegills. *Progr. Fish Cult.* 37(4):185-190.
- 12) Laal, A. K. and N. K. Thakur. 1977. Efficacy of grass carp (*Ctenopharyngodon idella*) in controlling *Potamogeton* infestation in a derelict pond in North Bihar. *Science and Culture* 43(9):391-392.
- 13) Shireman J. V. and M. J. Maceina. 1981. The utilization of grass carp, *Ctenopharyngodon idella* Val., for hydrilla control in Lake Baldwin, Florida. *J. Fish Biol.* 19:629-636.
- 14) Shireman, J. V., D. E. Colle and R. W. Rottmann. 1978. Size limits to predation on grass carp by largemouth bass. *T. Am. Fish. Soc.* 107(1):213-215.
- 15) Fowler, M. C. and T. O. Robson. 1978. The effects of the food preferences and stocking rates of grass carp (*Ctenopharyngodon idella* Val.) on mixed plant communities. *Aquat. Bot.* 5:261-276.
- 16) Baur, R. J., D. H. Buck and C. R. Rose. 1979. Production of age-0 largemouth bass, smallmouth bass, and bluegills in ponds stocked with grass carp. *Trans. Am. Fish. Soc.* 108:496-498.
- 17) Canfield, D. E. Jr., M. J. Maciena and J. V. Shireman. 1983. Effects of hydrilla and grass carp on water quality in a Florida lake. *Water Resour. B.* 19(5):773-778.
- 18) Leslie, A. J. Jr., L. E. Nall and J. M. Van Dyke. 1983. Effects of vegetation control by grass carp on selected water-quality variables in four Florida lakes. *Trans. Am. Fish. Soc.* 112:777-787.
- 19) Mitzner, L. 1978. Evaluation of biological control of nuisance aquatic vegetation by grass carp. *Trans. Am. Fish. Soc.* 107(1):135-145.
- 20) Shireman, J. V., D. E. Colle and M. J. Maceina. 1980. Grass carp growth rates in Lake Wales, Florida. *Aquaculture* 19:379-382.
- 21) Sutton, D. L. 1977. Grass carp (*Ctenopharyngodon idella* Val.) in North America. *Aquat. Bot.* 3:157-164.
- 22) Maceina, M. J. 1992. A simple regression model to assess environmental effects on fish growth. *J. Fish Biol.* 41:557-565.
- 23) Van Zon, J. C. J. 1977. Grass carp (*Ctenopharyngodon idella* Val.) in Europe. *Aquat. Bot.* 3:143-155.
- 24) Richard, D. I., J. W. Small Jr. and J. A. Osborne. 1984. Phytoplankton responses to reduction and elimination of submerged vegetation by herbicides and grass carp in four Florida lakes. *Aquat. Bot.* 20:307-319.
- 25) Lewis, W. M., R. C. Heidinger and R. O. Anderson. 1978. Summary and evaluation. *Trans. Am. Fish. Soc.* 107:223-224.
- 26) Kilgen, R. H. 1978. Growth of channel catfish and striped bass in small ponds stocked with grass carp and water hyacinths. *Trans. Am. Fish. Soc.* 107:176-180.
- 27) Forester, J. S. and J. W. Avault Jr. 1978. Effects of grass carp on freshwater red swamp crawfish in ponds. *Trans. Am. Fish. Soc.* 107:156-160.
- 28) Custer, P. E., F. D. Halverson, J. Malone and C. Von Chong. 1978. The white amur as a biological control agent of aquatic weeds in the Panama Canal. *Fisheries* 3(5):2-9.
- 29) Gasaway, R. D. 1977. Predation on introduced grass carp (*Ctenopharyngodon idella*) in a Florida lake. *Fla. Sci.* 40(2):167-173.
- 30) Gasaway, R. D. and T. F. Drda. 1978. Effects of grass carp introduction on macrophytic vegetation and chlorophyll content of phytoplankton in four Florida lakes. *Fla. Sci.* 41(2):101-109.
- 31) Beach, M. L. 1977. Grass carp for aquatic weed control, and its effect on water quality. *Fla. Sci.* 40:17.
- 32) Beach, M. L. 1977. Effects of grass carp on macro and microinvertebrates. *Fla. Sci.* 40:19.
- 33) Bain, M. B. 1989. Sterile grass carp may control hydrilla in Guntersville Reservoir. *Highlights of Agricultural Research, Alabama Agricultural Experimental Station* 36(3):5.
- 34) Tooby, T. E., J. Lucey and B. Stott. 1980. The tolerance of grass carp, *Ctenopharyngodon idella* Val., to aquatic herbicides. *J. Fish Biol.* 16:591-597.
- 35) Cassani, J. R. 1981. Feeding behaviour of underyearling hybrids of the grass carp, *Ctenopharyngodon idella* ♂ and the bighead, *Hypophthalmichthys nobilis* ♀ on selected species of aquatic plants. *J. Fish Biol.* 18:127-133.
- 36) Guillory, V. and R. D. Gasaway. 1978. Zoogeography of the grass carp in the United States. *Trans. Am. Fish. Soc.* 107:105-112.
- 37) Pflieger, W. L. 1978. Distribution and status of the grass carp (*Ctenopharyngodon idella*) in Missouri streams. *Trans. Am. Fish. Soc.* 107:113-118.
- 38) Stanley, J. G., W. W. Miley II and D. L. Sutton. 1978. Reproductive requirements and likelihood for naturalization of escaped grass carp in the United States. *Trans. Am. Fish. Soc.* 107:119-128.
- 39) Nixon, D. E. and R. L. Miller. 1978. Movements of grass carp, *Ctenopharyngodon idella*, in an open reservoir system as determined by underwater telemetry. *Trans. Am. Fish. Soc.* 107:146-148.
- 40) Colle, D. E., J. V. Shireman and R. W. Rottmann. 1978. Food selection by grass carp fingerlings in a vegetated pond. *Trans. Am. Fish. Soc.* 107:149-152.
- 41) Lewis, W. M. 1978. Observations on the grass carp in ponds containing fingerling channel catfish and hybrid sunfish. *Trans. Am. Fish. Soc.* 107:153-155.
- 42) Maceina, M. J., P. W. Bettoli, W. G. Klussmann, R. K. Betsill and R. L. Noble. 1991. Effect of aquatic macrophyte removal on recruitment and growth of black crappies and white crappies in Lake Conroe, Texas. *N. Am. J. Fish. Manage.* 11:556-563.
- 43) Cassani, J. R. 1995. Problems and prospects for grass carp as a management tool. *Am. Fish. Soc. Symp.* 15:407-412.
- 44) Fry, D. L. and J. A. Osborne. 1980. Zooplankton abundance and diversity in central Florida grass carp ponds. *Hydrobiologia* 68:145-155.
- 45) Del Fosse, E. S., D. L. Sutton and B. D. Perkins. 1976. Combination of the mottled waterhyacinth weevil and the white amur for biological control of waterhyacinth. *J. Aquat. Plant Manage.* 14:64-67.
- 46) Bailey, W. M. and R. L. Boyd. 1972. Some observations on the white amur in Arkansas. *Hyacinth Control J.* 10:20-22.
- 47) Michewicz, J. E., D. L. Sutton and R. D. Blackburn. 1972. Water quality of small enclosures stocked with white amur. *Hyacinth Control J.* 10:22-25.
- 48) Hestand, R. S. and C. C. Carter. 1978. Comparative effects of grass carp and selected herbicides on macrophyte and phytoplankton communities. *J. Aquat. Plant Manage.* 16:43-50.
- 49) Osborne, J. A. and N. M. Sassic. 1979. Biological control of *Hydrilla verticillata* Royle with grass carp. *J. Aquat. Plant Manage.* 17:45-48.
- 50) Ware, F. J. and R. D. Gasaway. 1978. Effects of grass carp on native fish populations in two Florida lakes, pp. 324-335. *In: Proceedings of the 30th Annual Conference of Southeastern Association of Fish and Wildlife Agencies.*
- 51) Eades, R. and E. Steinkoenig. 1995. An evaluation of aquatic vegetation control by triploid grass carp in Virginia ponds. *Am. Fish. Soc. Symp.* 15:437-442.
- 52) Greenfield, D. W. 1973. An evaluation of the advisability of the release of the grass carp, *Ctenopharyngodon idella*, into the natural waters of the United States. *Trans. Illinois Acad. Sci.* 66:47-53.
- 53) Rottmann, R. W. and R. O. Anderson. 1977. Limnological and ecological effects of grass carp in ponds, pp. 24-39. *In: Proceedings of the 30th Annual Conference of Southeastern Association of Fish and Wildlife Agencies.*

- 54) Miley, W. W. II, J. M. Van Dyke and D. M. Riley. 1979. The use of grass carp for biocontrol of aquatic weeds and their implication for natural resources and fisheries in Florida, pp. 159-175. *In: Proceedings of the Grass Carp Conference* (1978).
- 55) Klussmann, W. G., R. L. Noble, R. D. Martyn, W. J. Clark, R. K. Betsill, P. W. Bettoli, M. F. Cichra and J. M. Campbell. 1988. Control of aquatic macrophytes by grass carp in Lake Conroe, Texas, and the effects on the reservoir ecosystem. Texas Agricultural Experiment Station Miscellaneous Publication (1664). 61 pp.
- 56) Opuszynski, K. and J. V. Shireman. 1995. Herbivorous fishes, culture and use for weed management. CRC Press, Inc. 223 pp.
- 57) Van Zon, J. C. J. 1985. Conflicts of interest in the use of grass carp, pp. 399-403. *In: Proceedings of the VI International Symposium on Biological Control of Weeds* (1984).
- 58) Shireman, J. V., M. V. Hoyer, M. J. Maciena and D. E. Canfield Jr. 1985. The water quality and fishery of Lake Baldwin, Florida: 4 years after macrophyte removal by grass carp. Lake and Reservoir Management: Practical Applications, pp. 201-206. *In: Proceedings of the Fourth Annual Conference and International Symposium* (1984).
- 59) Chilton, E. W. II and S. M. Poarch. 1997. Distribution and movement behavior of radio-tagged grass carp in two Texas reservoirs. *Trans. Am. Fish. Soc.* 126:467-476.
- 60) Blackwell, B. G. and B. R. Murphy. 1996. Low-diversity triploid grass carp stockings for submersed vegetation control in small impoundments. *J. Freshw. Ecol.* 11 (4):475-484.
- 61) Ewel, K. C. and T. D. Fontaine III. 1976. Possible effects of the introduction of the white amur into Lake Conway, Florida, pp. 64-84. *In: Proceedings, Research Planning Conference on the Aquatic Plant Control Program* (1975).
- 62) Guillory, V., R. Land and R. Gasaway. 1977. Baseline data report-Lake Conway grass carp project, pp. 97-112. *In: Proceedings, Research Planning Conference on the Aquatic Plant Control Program* (1976).
- 63) Haller, W. T. and D. L. Sutton. 1976. Biocontrol of aquatic plant growth in earthen ponds by the white amur (*Ctenopharyngodon idella* Val), pp. 261-268. *In: Proceedings of the IV International Symposium on Biological Control of Weeds*.
- 64) Martyn, R. D., R. L. Noble, P. W. Bettoli and R. C. Maggio. 1986. Mapping aquatic weeds with aerial color infrared photography and evaluating their control by grass carp. *J. Aquat. Plant Manage.* 24:46-56.
- 65) Kasza, H., W. Krzanowski, E. Krzyzaneck and T. Kuflikowski. 1987. Bionotic changes in the Goczkalkowice Reservoir caused by the impact of grass carp (*Ctenopharyngodon idella* Val.) on macrophytes. *Bull. Pol. Acad. Sci.-Biol. Sci.* 35:189-197.
- 66) Pine, R. T. and L. W. J. Anderson. 1991. Effect of triploid grass carp on submersed aquatic plants in Northern California ponds. *California Fish and Game* 77(1):27-35.
- 67) Maceina, M. J., P. W. Bettoli, W. G. Klussman, R. K. Betsill and R. L. Noble. 1991. Effect of aquatic macrophyte removal on recruitment and growth of black crappies and white crappies in Lake Conroe, Texas. *N. Am. J. Fish. Manage.* 11:556-563.
- 68) Morrow, J. V. Jr. and J. P. Kirk. 1995. Age and growth of grass carp in Lake Guntersville, Alabama. *Proceedings of the Annual Conference of Southeastern Association of Fish and Wildlife Agencies* 49:187-194.
- 69) Morrow, J. V. Jr., J. P. Kirk and K. J. Killgore. 1997. Collection, age, growth, and population attributes of triploid grass carp stocked into the Santee-Cooper Reservoirs, South Carolina. *N. Am. J. Fish. Manage.* 17(1):38-43.
- 70) Osborne, J. A. 1985. A preliminary study of the efficacy of hybrid grass carp for hydrilla control. *J. Aquat. Plant Manage.* 23:16-20.
- 71) Pine, R. T., L. W. J. Anderson and S. S. O. Hung. 1990. Control of aquatic plants in static and flowing water by yearling triploid grass carp. *J. Aquat. Plant Manage.* 28:36-40.
- 72) Kilambi, R. V. and W. R. Robison. 1979. Effects of temperature and stocking density on food consumption and growth of grass carp *Ctenopharyngodon idella*, Val. *J. Fish Biol.* 15:337-342.
- 73) Forester, T. S. and J. M. Lawrence. 1978. Effects of grass carp and carp on populations of bluegill and largemouth bass in ponds. *Trans. Am. Fish. Soc.* 107:172-175.
- 74) Kirk, J. P. 1992. Efficacy of triploid grass carp in controlling nuisance aquatic vegetation in South Carolina farm ponds. *N. Am. J. Fish. Manage.* 12(3):581-584.
- 75) Lembi, C. A., B. G. Ritenour, E. M. Iverson and E. C. Forss. 1978. The effects of vegetation removal by grass carp on water chemistry and phytoplankton in Indiana ponds. *Trans. Am. Fish. Soc.* 107:161-171.
- 76) Prowse, G. A. 1971. Experimental criteria for studying grass carp feeding in relation to weed control. *Prog. Fish Cult.* 33(3):128-131.
- 77) Hill, K. R. 1986. Mortality and standing stocks of grass carp planted in two Iowa lakes. *N. Am. J. Fish. Manage.* 6:449-451.
- 78) Bailey, W. M. 1978. A comparison of fish populations before and after extensive grass carp stocking. *Trans. Am. Fish. Soc.* 107:181-206.
- 79) Venkatesh, B. and H. P. C. Shetty. 1978. Studies on the growth rate of the grass carp, *Ctenopharyngodon idella* (Valenciennes), fed on two aquatic weeds *ad libitum*. *Mysore J. Agr. Sci.* 12:622-628.
- 80) Avault, J. W. Jr. and A. Merkowsky. 1978. Evaluation of hybrid (common X grass) carp for weed control, pp. 184-193. *In: Symposium on Culture of Exotic Fishes*.
- 81) Krzywosz, T., W. Krzywosz and J. Radziej. 1980. The effect of grass carp, *Ctenopharyngodon idella* (Val.), on aquatic vegetation and ichthyofauna on Lake Dgal Wielki. *Ekol. Pol.-Pol. J. Ecol.* 28:433-450.
- 82) Tsuchiya, M. 1979. Natural reproduction of grass carp in the Tone River and their pond spawning, pp. 185-200. *In: Proceedings of the Grass Carp Conference* (1978).
- 83) Schoonbee, H. J., J. Vermaak and J. H. Swanepoel. 1985. Use of the Chinese grass carp, *Ctenopharyngodon idella*, in the control of the submerged water weed *Potamogeton pectinatus* in an inland lake in the Transvaal, South Africa, pp. 557-565. *In: Proceedings of the VI International Symposium on the Biological Control of Weeds*, (1984).
- 84) Mitchell, C. P. 1980. Control of water weeds by grass carp in two small lakes. *New Zeal. J. Mar. Fresh.* 14(4):381-390.
- 85) Rowe, D. K. 1984. Some effects of eutrophication and the removal of aquatic plants by grass carp (*Ctenopharyngodon idella*) on rainbow trout (*Salmo gairdnerii*) in Lake Parkinson, New Zealand. *New Zeal. J. Mar. Fresh.* 18:115-127.
- 86) Van Zon, J. C. J. 1984. Economic weed control with grass carp. *Trop. Pest Manage.* 30(2):179-185.
- 87) Lewis, W. M. 1974. The use of the white amur (*Ctenopharyngodon idella*) for control of aquatic weeds in connection with fish farming, p. 37. *In: Abstracts of the Meeting of the Weed Science Society of America*.
- 88) Terrell, J. W. 1974. Macrophyte control by the white amur in two catfish ponds, p. 37. *In: Abstracts of the Meeting of the Weed Science Society of America*.
- 89) Theriot, R. F. and D. R. Sanders Sr. 1974. Food preferences of yearling hybrid carp (*Ctenopharyngodon idella* × *Cyprinus carpio*), p. 37. *In: Abstracts of the Meeting of the Weed Science Society of America*.
- 90) Santha, C. R., W. E. Grant, W. H. Neill and R. K. Strawn. 1991. Biological control of aquatic vegetation using grass carp: simulation of alternative strategies. *Ecol. Model.* 59:229-245.
- 91) Tanner, C. C., R. D. S. Wells and C. P. Mitchell. 1990. Re-establishment of native macrophytes in Lake Parkinson following weed control by grass carp. *New Zeal. J. Mar. Fresh.* 24:181-186.
- 92) Mitchell, C. P. 1986. Effects of introduced grass carp on populations of two species of small native fishes in a small lake. *New Zeal. J. Mar. Fresh.* 20:219-230.
- 93) Van der Eijk, M. 1978. Notes on the experimental introduction of grass carp (*Ctenopharyngodon idella*) into the Netherlands, pp. 245-251. *In: Proceedings of the EWRS 5th Symposium on Aquatic Weeds*.
- 94) Mitzner, L. R. 1980. The use of the white amur in Iowa. *Proc. South. Weed Sci. Soc.* 33:204.
- 95) Pine, R. T. and L. W. J. Anderson. 1991. Plant preferences of triploid grass carp. *J. Aquat. Plant Manage.* 29:80-82.
- 96) Miley, W. W. II, J. M. Van Dyke, A. J. Leslie Jr., G. J. Kobylinski. 1980. Final report on weed control in four Florida lakes using grass carp. *Proc. South. Weed Sci. Soc.* 33:211-212.
- 97) Zweerde, W. V. D., B. J. Hoogers and J. C. J. V. Zon. 1978. Effects of grass carp on microflora and -fauna, macroflora, macroinvertebrates and chemical properties in the water, pp. 343-350. *In: Proceedings of the EWRS 5th Symposium on Aquatic Weeds*.
- 98) Venkatesh, B. and H. P. C. Shetty. 1978. Influence of the grass carp, *Ctenopharyngodon idella* (Valenciennes), on the physico-chemical and biological qualities of the fish pond, in relation to the feeds given. *Mysore J. Agr. Sci.* 12:605-617.
- 99) Riemens, R. G. 1978. Grass carp (*Ctenopharyngodon idella* Val.) as a tool in fishery management, pp. 351-358. *In: Proceedings of the EWRS 5th Symposium on Aquatic Weeds*.
- 100) Van Zon, J. C. J. 1980. Status of the use of grass carp, pp. 249-260. *In: Proceedings of V International Symposium on the Biological Control of Weeds*.

- 101) Mestrov, M. and V. Tavcar. 1973. Problems of recovery of Trakoscan Lake and application of grass carp. *Bull. Scientifique* 18(4-6):80-81.
- 102) Fowler, M. C. 1984. The present status of grass carp (*Ctenopharyngodon idella*) for the control of aquatic weeds in England and Wales, pp. 537-542. *In: Proceedings of VI International Symposium on the Biological Control of Weeds.*
- 103) Vinogradov, V. K. and Z. K. Zolotova. 1974. The influence of the grass carp on aquatic ecosystems. *Hydrobio.* 10(2):72-78.
- 104) Mitzner, L. 1979. Research and management application of grass carp in Iowa, pp. 31-48. *In: Proceedings of the Grass Carp Conference (1978).*
- 105) Shireman, J. V., D. E. Colle and R. G. Martin. 1979. Ecological study of Lake Wales, Florida after introduction of grass carp (*Ctenopharyngodon idella*), pp. 49-89. *In: Proceedings of the Grass Carp Conference (1978).*
- 106) Vogt, G. B. and H. A. Cordo. 1976. Recent South American field studies of prospective biocontrol agents of weeds, pp. 36-126. *In: Proceedings of the Research Planning Conf. on the Aquatic Plant Control Program (1975).*
- 107) Bailey, W. M. 1975. Operational experiences with the white amur in weed control programs, pp. 75-78. *In: Proceedings of the Symposium on Water Quality Management Through Biological Control.*
- 108) Beach, M. L., R. L. Lazor and A. P. Burkhalter. 1978. Some aspects of the environmental impact of the white amur (*Ctenopharyngodon idella* Val.) in Florida, and its use for aquatic weed control, pp. 269-289. *In: Proceedings of the IV International Symposium on the Biological Control of Weeds.*
- 109) Van Zon, J. C. J. 1979. The use of grass carp in comparison with other aquatic weed control methods, pp. 15-24. *In: Proceedings of the Grass Carp Conference (1978).*
- 110) Cassani, J. R., E. L. de la Vega and H. Allaire. 1995. An assessment of triploid grass carp stocking rates in small warmwater impoundments. *N. Am. J. Fish. Manage.* 15:400-407.
- 111) Shelton, W. L. and G. L. Jensen. 1979. Production of reproductively limited grass carp for biological control of aquatic weeds. *Water Resources Research Institute, Auburn Univ., Bulletin* 39. 173 pp.
- 112) Gasaway, R. D. and T. F. Drda. 1977. Effects of grass carp introduction on waterfowl habitat. *Trans. N. Am. Wildl. Nat. Res.* 42:73-85.
- 113) Willey, R. G., M. J. Dorskocil and C. A. Lembi. 1974. Potential of the white amur (*Ctenopharyngodon idella* Val.) as a biological control for aquatic weeds in Indiana, pp. 173-178. *In: Proc. Indiana Acad. Sci.*
- 114) Holdren, G. C., J. D. Mayfield and S. D. Porter. 1987. The effects of grass carp on water quality in McNeely Lake. *Lake Reserv. Manage.* 3:277-284.
- 115) Van, Zon, J. C. J., W. van der Zweerde and B. J. Hoogers. 1976. The grass-carp, its effects and side-effects, pp. 251-256. *In: Proceedings of IV International Symposium on Biological Control of Weeds.*
- 116) Miley, W. W. II, D. L. Sutton and J. G. Stanley. 1979. The role and impact of introduced grass carp (*Ctenopharyngodon idella* Val.) in the Union of Soviet Socialist Republics and several other European countries, pp. 177-183. *In: Proceedings of the Grass Carp Conference (1978).*
- 117) Henderson, S. 1979. Grass carp: the scientific and policy issues, pp. 25-29. *In: Proceedings of the Grass Carp Conference (1978).*
- 118) Stott, B. 1979. Grass carp research and public policy in England, pp. 147-158. *In: Proceedings of the Grass Carp Conference (1978).*
- 119) Venter, A. J. A. and H. J. Schoonbee. 1991. The use of triploid carp, *Ctenopharyngodon idella* (Val.), in the control of submerged aquatic weeds in the Florida lake, Roodepoort, Transvaal. *Water S. A.* 17(4):321-326.
- 120) Sutton, D. L., V. V. Van Diver Jr., R. S. Hestand and W. W. Miley II. 1979. Use of the grass carp for control of hydrilla in small ponds, pp. 91-102. *In: Proceedings of the Grass Carp Conference (1978).*
- 121) Mitzner, L. 1976. Biological control of nuisance aquatic macrophytes by white amur, pp. 105-106. *In: Proceedings of the North-central Weed Control Conference.*
- 122) Theriot, R. F. and J. L. Decell. 1978. Large-scale operations management test of the use of the white amur to control aquatic plants, pp. 220-229. *In: Symposium on Culture of Exotic Fishes.*
- 123) Fedorenko, A. Y. and F. J. Fraser. 1978. A review of the biology of grass carp (*Ctenopharyngodon idella* Val.) and its evaluation as a potential weed control agent in British Columbia. *Fisheries and Marine Service Technical Report No. 786.* 15p.
- 124) Ewel, K. C. and T. D. Fontaine III. 1976. Proposed relationships between white amur and the aquatic ecosystem at Lake Conway, Florida, pp. 159-176. *In: Proceedings of Research and Planning Conference on the Aquatic Plant Control Program.*
- 125) Wiley, M. J., P. P. Tazik and S. T. Sobaski. 1987. Controlling aquatic vegetation with triploid grass carp. *Circular 57.* Illinois Natural History Survey, Champaign. 16 p.
- 126) Cross, D. G. 1969. Aquatic weed control using grass carp. *J. Fish Biol.* 1:27-30.
- 127) Kilambi, R. V. and A. Zdinak. 1980. Food preference and growth of grass carp, *Ctenopharyngodon idella*, and hybrid carp, *C. idella* female x *Aristichthys nobilis* male, pp. 281-286. *In: Proceedings of V International Symposium on Biological Control of Weeds.*
- 128) King, D. R. and G. S. Hunt. 1967. Effect of carp on vegetation in a Lake Erie marsh. *J. Wildlife Manage.* 31(1):181-188.
- 129) Shelton, W. L., R. O. Smitherman and G. L. Jensen. 1981. Density related growth of grass carp, *Ctenopharyngodon idella* Val. in managed small impoundments in Alabama. *J. Fish Biol.* 18:45-51.
- 130) Kilgen, R. H. and R. O. Smitherman. 1971. Food habits of the white amur stocked in ponds alone and in combination with other species. *Prog. Fish Cult.* 33(3):123-127.
- 131) Duthu, G. S. and R. H. Kilgen. 1975. Aquarium studies on the selectivity of 16 aquatic plants as food by fingerling hybrids of the cross between *Ctenopharyngodon idella* and *Cyprinus carpio*. *J. Fish Biol.* 7:203-208.
- 132) Stott, B., D. G. Cross, R. E. Iszard and T. O. Robson. 1971. Recent work on grass carp in the United Kingdom from the standpoint of its economics in controlling submerged aquatic plants, pp. 105-116. *In: Proceedings of European Weed Research Society, 3rd International Symposium on Aquatic Weeds.*
- 133) Bain, M. B., D. H. Webb, M. D. Trangedal and L. N. Mangum. 1990. Movements and habitat use by grass carp in a large mainstream reservoir. *Trans. Am. Fish. Soc.* 119(3):553-561.
- 134) Bonar, S. A., G. L. Thomas, S. L. Thiesfeld, G. B. Pauley and T. B. Stables. 1993. Effect of triploid grass carp on the aquatic macrophyte community of Devils Lake, Oregon. *N. Am. J. Fish. Manage.* 13(4):757-765.
- 135) Clapp, D. F., R. S. Hestand III, B. Z. Thompson and L. L. Connor. 1993. Movement of triploid grass carp in large Florida lakes. *N. Am. J. Fish. Manage.* 13(4):746-756.
- 136) Beyers, D. W. and C. A. Carlson. 1993. Movement and habitat use of triploid and diploid grass carp in a Colorado irrigation canal. *N. Am. J. Fish. Manage.* 13(1):141-150.
- 137) Maciena, M. J., M. F. Cichra, R. K. Betsill and P. W. Bettoli. 1992. Limnological changes in a large reservoir following vegetation removal by grass carp. *J. Freshwater Ecol.* 7(1):81-95.
- 138) Cui, Y. B., S. L. Chen and S. M. Wang. 1994. Effect of ration size on the growth and energy budget of the grass carp. *Aquaculture* 123(1-2):95-107.
- 139) Bain, M. B. 1993. Assessing impacts of introduced aquatic species-grass carp in large systems. *Environ. Manage.* 17(2):211-224.
- 140) Cui, Y. B., S. L. Chen, S. M. Wang and X. F. Liu. 1993. Laboratory observations on the circadian feeding patterns in the grass carp fed 3 different diets. *Aquaculture* 113(1-2):57-64.
- 141) Bettoli, P. W., M. J. Maciena, R. L. Noble and R. K. Betsill. 1993. Response of a reservoir fish community to aquatic vegetation removal. *N. Am. J. Fish. Manage.* 13(1):110-124.
- 142) Rottmann, R. W. 1976. Limnological and ecological effects of grass carp in ponds. *M.S. thesis.* 62 pp.
- 143) Sills, J. B. 1970. A review of herbivorous fish for weed control. *Prog. Fish Cult.* 32(3):158-161.
- 144) Hawker, J. L. 1976. Whither the grass carp? *Conservationist* 37(5):16-19.
- 145) Kilambi, R. V. and M. L. Galloway. 1985. Temperature preference and tolerance of hybrid carp (female grass carp, *Ctenopharyngodon idella* x male bighead, *Aristichthys nobilis*). *Environ. Biol. Fish.* 12(4):309-314.
- 146) Kilambi, R. V. and A. Zdinak. 1982. Food intake and growth of hybrid carp (female grass carp, *Ctenopharyngodon idella* x male bighead, *Aristichthys (Hypophthalmichthys) nobilis*) fed on zooplankton and *Chara*. *J. Fish Biol.* 21:63-67.
- 147) Kilambi, R. V. and A. Zdinak. 1981. Comparison of early developmental stages and adults of grass carp, *Ctenopharyngodon idella*, and hybrid carp (female grass carp x male bighead *Aristichthys nobilis*). *J. Fish Biol.* 19:457-465.

- 148) Januszko, M. 1974. The effect of three species of phytophagous fish on algae development. *Pol. Arch. Hydrobiol.* 21:431-454.
- 149) Santha, C. R., R. D. Martyn, W. H. Neill and K. Strawn. 1994. Control of submersed weeds by grass carp in waterlily production ponds. *J. Aquat. Plant Manage.* 32:29-33.
- 150) Van Donk, E. and A. Otte. 1996. Effects of grazing by fish and waterfowl on the biomass and species composition of submerged macrophytes. *Hydrobiologia* 340:285-290.
- 151) Durocher, P. P., W. C. Provine and J. E. Kraai. 1984. Relationship between abundance of largemouth bass and submerged vegetation in Texas Reservoirs. *N. Am. J. Fish. Manage.* 4:84-88.
- 152) Brown, D. J. and T. G. Coon. 1991. Grass carp larvae in the lower Mississippi River and its tributaries. *N. Am. J. Fish. Manage.* 11:62-66.
- 153) Zimpfer, S. P., C. F. Bryan and C. H. Pennington. 1987. Factors associated with the dynamics of grass carp larvae in the lower Mississippi River valley. *Am. Fish. Soc. Symp.* 2:102-108.
- 154) Bonar, S. A., B. Bolding and M. Divens. 2002. Effects of triploid grass carp on aquatic plants, water quality, and public satisfaction in Washington State. *N. Am. J. Fish. Manage.* 22(1):96-105.
- 155) Hanlon, S. G., M. V. Hoyer, C. E. Cichra and D. E. Canfield. 2000. Evaluation of macrophyte control in 38 Florida lakes using triploid grass carp. *J. Aquat. Plant Manage.* 38:48-54.
- 156) Hargrave, C. W. and K. B. Gido. 2004. Evidence of reproduction by exotic grass carp in the Red and Washita rivers, Oklahoma. *Southwest. Nat.* 49(1):89-93.
- 157) Killgore, K. J., J. P. Kirk and J. W. Foltz. 1998. Response of littoral fishes in Upper Lake Marion, South Carolina following Hydrilla control by triploid grass carp. *J. Aquat. Plant Manage.* 36:82-87.
- 158) Kirk, J. P., J. V. Morrow, K. J. Killgore, S. J. De Kozlowski and J. W. Preacher. 2000. Population response of triploid grass carp to declining levels of Hydrilla in the Santee Cooper reservoirs, South Carolina. *J. Aquat. Plant Manage.* 38:14-17.
- 159) Kirk, J. P. and R. C. Socha. 2003. Longevity and persistence of triploid grass carp stocked into the Santee Cooper reservoirs of South Carolina. *J. Aquat. Plant Manage.* 41:90-92.
- 160) Kirkagac, M. and N. Demir. 2004. The effects of grass carp on aquatic plants, plankton and benthos in ponds. *J. Aquat. Plant Manage.* 42:32-39.
- 161) Pipalova, I. 2002. Initial impact of low stocking density of grass carp on aquatic macrophytes. *Aquat. Bot.* 73(1):9-18.
- 162) Kirk, J. P., K. J. Killgore, J. V. Morrow Jr., S. D. Lamprecht and D. W. Cooke. 2001. Movements of triploid grass carp in the Cooper River, South Carolina. *J. Aquat. Plant Manage.* 39:59-62.
- 163) Bonar, S. A., H. S. Sehgal, G. B. Pauley and G. L. Thomas. 1990. Relationship between the chemical composition of aquatic macrophytes and their consumption by grass carp, *Ctenopharyngodon idella*. *J. Fish Biol.* 36:149-157.
- 164) Wiley, M. J., S. M. Pescitelli and L. D. Wike. 1986. The relationship between feeding preferences and consumption rates in grass carp and grass carp x bighead carp hybrids. *J. Fish Biol.* 29:507-514.
- 165) Murphy, J. E., K. B. Beckmen, J. K. Johnson, R. B. Cope, T. Lawmaster and V. R. Beasley. 2002. Toxic and feeding deterrent effects of native aquatic macrophytes on exotic grass carp (*Ctenopharyngodon idella*). *Ecotoxicology* 11:243-254.
- 166) Vincent, J. F. V. and F. A. Sibbing. 1992. How the grass carp (*Ctenopharyngodon idella*) chooses and chews its food - some clues. *J. Zool.* 226:435-444.